

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 237 362 B2

(12)

## NEW EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention  
of the opposition decision:  
21.10.1998 Bulletin 1998/43

(51) Int Cl.<sup>6</sup>: C12Q 1/68, C12Q 1/70

(45) Mention of the grant of the patent:  
11.03.1992 Bulletin 1992/11

(21) Application number: 87302196.8

(22) Date of filing: 13.03.1987

(54) **Process for detecting specific nucleotide variations and genetic polymorphisms present in nucleic acids and kits therefor**

Verfahren und Ausstattung zur Bestimmung von spezifischen Nukleotid-Variationen und genetischen Polymorphismen in Nukleinsäuren

Procédé et nécessaire pour la détection des variations de nucléotides spécifiques et de polymorphismes génétiques dans les acides nucléiques

(84) Designated Contracting States:  
AT BE CH DE FR GB IT LI LU NL SE

(30) Priority: 13.03.1986 US 839331  
22.08.1986 US 899344

(43) Date of publication of application:  
16.09.1987 Bulletin 1987/38

(60) Divisional application: 91113105.0 / 0 459 532  
91113236.3 / 0 459 533

(73) Proprietor: F. HOFFMANN-LA ROCHE AG  
4002 Basel (CH)

(72) Inventors:

- Erlich, Henry Anthony  
Oakland California 94602 (US)
- Saiki, Randall Keichi  
Richmond California 94805 (US)
- Horn, Glenn Thomas  
Emeryville California 94608 (US)
- Mullis, Kary Banks  
La Jolla California 92037 (US)

(74) Representative: VOSSIUS & PARTNER  
Postfach 86 07 67  
81634 München (DE)

(56) References cited:

EP-A- 0 111 340	EP-A- 0 123 513
EP-A- 0 130 515	EP-A- 0 164 054
EP-A- 0 194 545	WO-A-84/04332
US-A- 3 909 363	US-A- 4 521 509

- CHEMICAL ABSTRACTS, vol. 103, 1985, page 186, abstract no. 117310n, Columbus, Ohio, US; J.W. GAUBATZ et al.: "Displacement synthesis of globin complementary DNA: evidence for sequence amplification", & BIOCHIM. BIOPHYS. ACTA 1985, 825(2), 175-87
- CHEMICAL ABSTRACTS, vol. 104, 1986, page 163, abstract no. 62996d, Columbus, Ohio, US; D.J. CUMMINGS et al.: "Excision-amplification of mitochondrial DNA during senescence in Podospora anserina. DNA sequence analysis of three unique 'plasmids'", & J. MOL. BIOL. 1985, 185(4), 659-80
- CHEMICAL ABSTRACTS, vol. 99, 1983, page 156, abstract no. 153092a, Columbus, Ohio, US; A.V. GUDKOV et al.: "Gene amplification in mammalian somatic cells resistant to colchicine. Part VI. Analysis of amplified DNA sequences by restriction endonuclease digestion", & GENETIKA (MOSCOW) 1983, 19(7), 1045-53
- Methods in Enzymology; vol.100, 1983, p.266-85
- Science; vol.230, 1985, p.1350-54
- Nucl. Acids Res.; vol.7, no.6, 1979, p.1541-52
- Proc. Natl. Acad. Sci. USA; vol.80, 1983, p.4045-49

## Remarks:

Divisional application 91113105.0 filed on 13/03/87.

EP 0 237 362 B2

## Description

This invention relates to a process for detecting nucleotide variations, mutations and polymorphisms by amplifying nucleic acid sequences suspected of containing such mutations or polymorphisms and detecting them with sequence-specific oligonucleotides in a dot blot format.

In recent years, the molecular basis of a number of human genetic diseases has been elucidated by the application of recombinant DNA technology. In particular, the detection of specific polymorphic restriction sites in human genomic DNA associated with genetic disease, such as sickle-cell anemia, has provided clinically valuable information for pre-natal diagnosis. In these studies, the presence or absence of a specific site is revealed by restriction fragment length polymorphism (RFLP) analysis, a method in which variation in the size of a specific genomic restriction fragment is detected by Southern blotting and hybridization of the immobilized genomic DNA with a labeled probe. RFLP analysis has proved useful in the direct detection of polymorphic sites that contain the mutation conferring the disease phenotype (e.g., MstII and sickle-cell anemia) as well as in linkage studies where a particular allelic restriction site is linked to a disease locus within a family but not necessarily in the general population. See, for example, Kan and Dozy, PNAS (USA), 75, 5631 (1978), and Rubin and Kan, Lancet, 1985-I, 75 (1985). See also Geever et al., PNAS (USA), 78, 5081 (1981) and Wilson et al., PNAS (USA), 79 3628 (1982).

In a second method, called "oligomer restriction", a synthetic end-labeled oligonucleotide probe is annealed in solution to the target genomic DNA sequence and a restriction enzyme is added to cleave any hybrids formed. This method, the specificity of which depends on the ability of a base pair mismatch within the restriction site to abolish or inhibit cleavage, is described more fully by Saiki et al., Biotechnology, 3, 1008-1012 (1985). In addition, the sensitivity of this technique may be enhanced by utilizing a polymerase chain reaction procedure wherein the sample is first subjected to treatment with specific primers, polymerase and nucleotides to amplify the signal for subsequent detection. This is described more fully by Saiki et al., Science, 230, 1350-1353 (1985).

A third method for detecting allelic variations which is independent of restriction site polymorphism utilizes sequence-specific synthetic oligonucleotide probes. See Conner et al., PNAS (USA), 80, 78 (1983). This latter technique has been applied to the prenatal diagnosis of  $\alpha$ 1-antitrypsin deficiency (Kidd et al., Nature, 304, 230 (1983)) and  $\beta$ -thalassemia (Pirastu et al., N. Engl. J. Med., 309, 284 (1983)). In addition, the technique has been applied to study the polymorphism of HLA-DR $\beta$  using Southern blotting (Angelini et al., PNAS (USA), 83, 4489-4493 (1986)).

The basis for this procedure is that under appropriate hybridization conditions a short oligonucleotide probe of at least 19 bases (19-mer) will anneal only to those sequences to which it is perfectly matched, a single base pair mismatch being sufficiently destabilizing to prevent hybridization. The distinction between the allelic variants is based on the thermal stability of the duplex formed between the genomic DNA and the oligonucleotide (19-mer) probe.

In addition, methods for detecting base pair mismatches in double-stranded RNA and RNA:DNA heteroduplexes have been described using pancreatic ribonuclease (RNase A) to cleave the heteroduplexes. Winter et al., PNAS (USA), 82:7575-7579 (1985) and Myers et al., Science, 230:1242-1246 (1985). The principal deficiency of this method is its inability to recognize all types of base pair mismatches.

Both the RFLP and oligonucleotide probe stability methods are relatively complex procedures, requiring restriction enzyme digestion, gel-fractionation of the genomic DNA, denaturation of the DNA, immobilization of the DNA either by transfer to a filter membrane or dessication of the gel itself, and hybridization of a labeled probe to the electrophoretically resolved array of immobilized genomic restriction fragments. These steps are necessary, for the oligonucleotide probe stability method, due to the complexity of human genomic DNA. Restriction and electrophoresis are necessary to separate the target sequence ("signal") from the rest of the genome ("noise"), and hybridization in the gel (instead of filter transfer) is necessary to retain as much target sequence as possible. Even then, detection of a signal in a 10  $\mu$ g sample using a high specific activity kinased probe requires an autoradiographic exposure of four days.

In addition, the approach of Conner et al. requires at least a 19-mer probe for reasons of specificity (a shorter probe would hybridize to more genomic fragments), as well as possibly sensitivity. Shorter probes (e.g., 16-mers), however, would show more sequence-specific discrimination because a single mismatch would be more destabilizing.

There is a need in the art for a simplified method with improved sensitivity and specificity for directly detecting single-base differences in nucleic acids such as genomic DNA.

Accordingly, the present invention provides a method for detecting single or multiple nucleotide variations in nucleic acid sequence from any source, for use in detecting any type of disease or condition. The method herein directly detects the sequence variation, eliminating the need for restriction digestion, electrophoresis, and gel manipulations otherwise required. In addition, the method herein provides for improved specificity and sensitivity of the probe; an interpretable signal can be obtained with a 0.04  $\mu$ g sample in six hours. Thirdly, if the amount of sample spotted on a membrane is increased to 0.1-0.5  $\mu$ g, non-isotopically labeled oligonucleotides may be utilized rather than the radioactive probes used in previous methods. Finally, the process herein is applicable to use of sequence-specific oligonucleotides less than 19-mers in size, thus allowing use of more discriminatory sequence-specific oligonucleotides.

Specifically, the present invention provides for a process for detecting the presence or absence of at least one

nucleotide variation in sequence in one or more nucleic acids contained in a sample, which process comprises:

- (a) treating the sample, together or sequentially, with four different nucleotide triphosphates, an agent for polymerization of the nucleotide triphosphates, and one oligonucleotide primer for each strand of each nucleic acid suspected of containing said variation under hybridizing conditions, such that for each nucleic acid strand containing each different variation to be detected, an extension product of each primer is synthesized which is complementary to each nucleic acid strand, wherein said primer or primers are selected so as to be substantially complementary to each nucleic acid strand containing each different variation, such that the extension product synthesized from one primer, when it is separated from its complement, can serve as a template for synthesis of the extension product of the other primer;
- (b) treating the sample under denaturing conditions to separate the primer extension products from their templates if the variation(s) to be detected are present;
- (c) treating the sample, together or sequentially, with said four nucleotide triphosphates, an agent for polymerization of the nucleotide triphosphates, and oligonucleotide primers such that a primer extension product is synthesized using each of the single strands produced in step (b) as a template, wherein steps (b) and (c) are repeated a sufficient number of times to result in detectable amplification of the nucleic acid containing the sequence variation (s), if present;
- (d) affixing the product of step (c) to a membrane;
- (e) treating the membrane under hybridization conditions with a labeled sequence-specific oligonucleotide probe capable of hybridizing with the amplified nucleic acid sequence only if the sequence of the probe is complementary to a region of the amplified sequence; and
- (f) detecting whether the probe has hybridized to an amplified sequence in the nucleic acid sample.

Steps (b) and (c) are preferably repeated at least five times, and more preferably 15-30 times if the sample contains human genomic DNA. If the sample comprises cells, preferably they are heated before step (a) to expose the nucleic acids therein to the reagents. This step avoids purification of the nucleic acids prior to reagent addition.

In a variation of this process, the primer(s) and/or nucleotide triphosphates are labeled so that the resulting amplified sequence is labeled. The labeled primer(s) and/or nucleotide triphosphate(s) can be present in the reaction mixture initially or added during a later cycle. The sequence-specific oligonucleotide (unlabeled) is affixed to a membrane and treated under hybridization conditions with the labeled amplification product so that hybridization will occur only if the membrane-bound sequence is present in the amplification product.

In another embodiment, the invention herein relates to a kit for detecting the presence of a specific nucleotide sequence in a nucleic acid in a sample, which kit comprises, in packaged form, a multicontainer unit having:

- (a) two oligonucleotide primers for said nucleic acid, which primers are selected such that an extension product synthesized from one primer, when separated from its complement, can serve as a template for the synthesis of the extension product of the other primer so as to exponentially amplify said nucleic acid;
- (b) an agent for polymerization;
- (c) four different nucleoside triphosphates; and either:
- (d) a sequence specific oligonucleotide probe capable of hybridizing with the amplified nucleic acid spotted directly on a membrane only if the sequence of the probe is complementary to a region of the amplified nucleic acid; and
- (e) a membrane to which amplified nucleic acid can directly bind for subsequent detection by probe hybridization; or
- (d) a sequence specific oligonucleotide probe attached to a membrane, wherein said probe is capable of hybridizing with the amplified nucleic acid only if the sequence of the probe is complementary to a region of the amplified nucleic acid.

Regarding genetic diseases, while RFLP requires a polymorphic restriction site to be associated with the disease, sequence-specific oligonucleotides directly detect the genetic lesion and are generally more useful for the analysis of such genetic diseases as hemoglobin C disease,  $\alpha$ 1-antitrypsin and  $\beta$ -thalassemia which result from single or multiple base mutations. In addition, the oligonucleotides can be used to distinguish between genetic variants which represent different alleles (e.g., HLA typing), indicating the feasibility of a sequence specific oligonucleotide-based HLA typing kit.

The term "nucleotide variation in sequence" refers to any single or multiple nucleotide substitutions, deletions or insertions. These nucleotide variations may be mutant or polymorphic allele variations. Therefore, the process herein can detect single nucleotide changes in nucleic acids such as occur in  $\beta$ -globin genetic diseases caused by single-base mutations, additions and deletions (some  $\beta$ -thalassemias, sickle cell anemia, hemoglobin C disease, etc.), as well as multiple-base variations such as are involved with  $\alpha$ -thalassemia or some  $\beta$ -thalassemias. The process can also detect polymorphisms, which are not necessarily associated with a disease, but are merely a condition in which two or more different nucleotide sequences (whether having substituted, deleted or inserted nucleotide base pairs)

can exist at a particular site in the nucleic acid in the population, as with HLA regions of the human genome and random polymorphisms such as in mitochondrial DNA. The polymorphic sequence-specific oligonucleotide probes described in detail hereinafter may be used as genetic markers linked to a disease such as insulin-dependent diabetes or in forensic applications. If the nucleic acid is double-stranded, the nucleotide variation in sequence becomes a base pair variation in sequence.

The term "oligonucleotide" as used herein is defined as a molecule comprised of two or more deoxyribonucleotides or ribonucleotides, preferably more than three. Its exact size will depend on many factors, which in turn depend on the ultimate function or use of the oligonucleotide. The oligonucleotide may be derived synthetically or by cloning.

The term "primer as used herein refers to an oligonucleotide, whether occurring naturally as in a purified restriction digest or produced synthetically, which is capable of acting as a point of initiation of synthesis when placed under conditions in which synthesis of a primer extension product which is complementary to a nucleic acid strand is induced, i.e., in the presence of four different nucleotide triphosphates and an agent for polymerization such as DNA polymerase in an appropriate buffer ("buffer" includes pH, ionic strength, cofactors, etc.) and at a suitable temperature.

The primer is preferably single stranded for maximum efficiency in amplification, but may alternatively be double stranded. If double stranded, the primer is first treated to separate its strands before being used to prepare extension products. Preferably, the primer is an oligodeoxyribonucleotide. The primer must be sufficiently long to prime the synthesis of extension products in the presence of the agent for polymerization. The exact lengths of the primers will depend on many factors, including temperature and source of primer and use of the method. For example, depending on the complexity of the target sequence, the oligonucleotide primer typically contains 15-25 nucleotides, although it may contain more or fewer nucleotides. Short primer molecules generally require lower temperatures to form sufficiently stable hybrid complexes with the template.

The primers herein are selected to be "substantially" complementary to the different strands of each specific sequence to be amplified. This means that the primers must be sufficiently complementary to hybridize with their respective strands. Therefore, the primer sequence need not reflect the exact sequence of the template. For example, a non-complementary nucleotide fragment may be attached to the 5' end of the primer, with the remainder of the primer sequence being complementary to the strand. Typically, the primers have exact complementarity to obtain the best detection results.

The term "sequence-specific oligonucleotides" refers to oligonucleotides which will hybridize to specific sequences, whether or not contained on alleles which sequences span the nucleotide variation being detected and are specific for the sequence variation being detected. Depending on the sequences being analyzed, one or more sequence-specific oligonucleotides may be employed for each sequence, as described further hereinbelow.

As used herein, the term "thermostable enzyme" refers to an enzyme which is stable to heat and is heat resistant and catalyzes (facilitates) combination of the nucleotides in the proper manner to form the primer extension products which are complementary to each nucleic acid strand. Generally, the synthesis will be initiated at the 3' end of each primer and will proceed in the 5' direction along the template strand, until synthesis terminates, producing molecules of different lengths. There may be thermostable enzymes, however, which initiate synthesis at the 5' end and proceed in the other direction, using the same process as described above. A purified thermostable enzyme is described more fully in Example VIII hereinbelow.

The present invention is directed to a process for amplifying any one or more specific nucleic acid sequences (as defined herein to contain one or more nucleotide variations) suspected of being in one or more nucleic acids.

In general, the present process involves a chain reaction for producing, in exponential quantities relative to the number of reaction steps involved, at least one specific nucleic acid sequence given (a) that the ends of the required sequence are known in sufficient detail that oligonucleotides can be synthesized which will hybridize to them, and (b) that a small amount of the sequence is available to initiate the chain reaction. The product of the chain reaction will be a discrete nucleic acid duplex with termini corresponding to the ends of the specific primers employed.

Any nucleic acid, in purified or non purified form, can be utilized as the starting nucleic acid or acids, provided it is suspected of containing the sequence being detected. Thus, the process may employ, for example, DNA or RNA, including messenger RNA, which DNA or RNA may be single stranded or double stranded. In addition, a DNA-RNA hybrid which contains one strand of each may be utilized. A mixture of any of these nucleic acids may also be employed, or the nucleic acids produced from a previous amplification reaction herein using the same or different primers may be so utilized. The specific nucleic acid sequence to be amplified may be only a fraction of a larger molecule or can be present initially as a discrete molecule, so that the specific sequence constitutes the entire nucleic acid.

It is not necessary that the sequence to be amplified be present initially in a pure form; it may be a minor fraction of a complex mixture, such as a portion of the  $\beta$ -globin gene contained in whole human DNA, or portion of nucleic acid sequence due to a particular microorganism which organism might constitute only a very minor fraction of a particular biological sample. The starting nucleic acid may contain more than one desired specific nucleic acid sequence which may be the same or different. Therefore, the present process is useful not only for producing large amounts of one specific nucleic acid sequence, but also for amplifying simultaneously more than one different specific nucleic acid

sequence located on the same or different nucleic acid molecules if more than one of the base pair variations in sequence is present.

The nucleic acid or acids may be obtained from any source, for example, from plasmids such as pBR322 from cloned DNA or RNA, or from natural DNA or RNA from any source, including bacteria, yeast, viruses, organelles, and higher organisms such as plants or animals. DNA or RNA may be extracted from blood, tissue material such as chorionic villi or amniotic cells by a variety of techniques such as that described by maniatitis et al., Molecular Cloning (1982), 280-281.

The cells may be directly used without purification of the nucleic acid if they are suspended in hypotonic buffer and heated to about 90-100°C, until cell lysis and dispersion of intracellular components occur generally about 1 to 15 minutes. After the heating step the amplification reagents may be added directly to the lysed cells. This direct cell detection method may be used on peripheral blood lymphocytes and amniocytes.

Any specific nucleic acid sequence can be amplified by the present process. It is only necessary that a sufficient number of bases at both ends of the sequence be known in sufficient detail so that two oligonucleotide primers can be prepared which will hybridize to different strands of the desired sequence and at relative positions along the sequence such that an extension product synthesized from one primer, when it is separated from its template (complement), can serve as a template for extension of the other primer into a nucleic acid of defined length. The greater the knowledge about the bases at both ends of the sequence, the greater can be the specificity of the primers for the target nucleic acid sequence, and thus the greater the efficiency of the process.

It will be understood that the word "primer" as used hereinafter may refer to more than one primer, particularly in the case where there is some ambiguity in the information regarding the terminal sequence(s) of the fragment to be amplified. For instance, in the case where a nucleic acid sequence is inferred from protein sequence information, a collection of primers containing sequences representing all possible codon variations based on degeneracy of the genetic code will be used for each strand. One primer from this collection will be homologous with the end of the desired sequence to be amplified.

The oligonucleotide primers may be prepared using any suitable method, such as, for example, the organic synthesis of a nucleic acid from nucleoside derivatives. This synthesis may be performed in solution or on a solid support. One type of organic synthesis is the phosphotriester method, which has been utilized to prepare gene fragments or short genes. In the phosphotriester method, oligonucleotides are prepared that can then be joined together to form longer nucleic acids. For a description of this method, see Narang, S. A., et al., Meth. Enzymol., 68, 90 (1979) and U. S. Patent No. 4,356,270. The patent describes the synthesis and cloning of the somatostatin gene.

A second type of organic synthesis is the phosphodiester method, which has been utilized to prepare a tRNA gene. See Brown, E. L., et al., Meth. Enzymol., 68, 109 (1979) for a description of this method. As in the phosphotriester method, this phosphodiester method involves synthesis of oligonucleotides that are subsequently joined together to form the desired nucleic acid.

Automated embodiments of these methods may also be employed. In one such automated embodiment, di ethyl phosphoramidites are used as starting materials and may be synthesized as described by Beaucage et al., Tetrahedron Letters (1981), 22:1859-1862. One method for synthesizing oligonucleotides on a modified solid support is described in U.S. Patent No. 4,458,066. It is also possible to use a primer which has been isolated from a biological source (such as a restriction endonuclease digest).

The specific nucleic acid sequence is produced by using the nucleic acid containing that sequence as a template. If the nucleic acid contains two strands, it is necessary to separate the strands of the nucleic acid before it can be used as the template, either as a separate step or simultaneously with the synthesis of the primer extension products. This strand separation can be accomplished by any suitable denaturing method including physical, chemical or enzymatic means. One physical method of separating the strands of the nucleic acid involves heating the nucleic acid until it is completely (>99%) denatured. Typical heat denaturation may involve temperatures ranging from about 80 to 105°C for times ranging from about 1 to 10 minutes. Strand separation may also be induced by an enzyme from the class of enzymes known as helicases or the enzyme RecA, which has helicase activity and in the presence of riboATP is known to denature DNA. The reaction conditions suitable for separating the strands of nucleic acids with helicases are described by Kuhn Hoffmann-Berling, CSH-Quantitative Biology, 43:63 (1978), and techniques for using RecA are reviewed in C. Radding, Ann. Rev. Genetics, 16:405-37 (1982).

If the original nucleic acid containing the sequence variation to be amplified is single stranded, its complement is synthesized by adding one or two oligonucleotide primers thereto. If an appropriate single primer is added, a primer extension product is synthesized in the presence of the primer, an agent for polymerization, and the four nucleotide triphosphates described below. The product will be partially complementary to the single-stranded nucleic acid and will hybridize with the nucleic acid strand to form a duplex of unequal length strands that may then be separated into single strands as described above to produce two single separated complementary strands. Alternatively, two appropriate primers may be added to the single-stranded nucleic acid and the reaction carried out.

If the original nucleic acid constitutes the entire sequence variation to be amplified, the primer extension product

(s) produced will be completely complementary to the strands of the original nucleic acid and will hybridize therewith to form a duplex of equal length strands to be separated into single-stranded molecules.

When the complementary strands of the nucleic acid or acids are separated, whether the nucleic acid was originally double or single stranded, the strands are ready to be used as a template for the synthesis of additional nucleic acid strands. This synthesis can be performed using any suitable method. Generally it occurs in a buffered aqueous solution, preferably at a pH of 7-9, most preferably about 8. Preferably, a molar excess (for cloned nucleic acid, usually about 1000:1 primer:template, and for genomic nucleic acid, usually about  $10^6$ :1 primer:template) of the two oligonucleotide primers is added to the buffer containing the separated template strands. It is understood, however, that the amount of complementary strand may not be known if the process herein is used for diagnostic applications, so that the amount of primer relative to the amount of complementary strand cannot be determined with certainty. As a practical matter, however, the amount of primer added will generally be in molar excess over the amount of complementary strand (template) when the sequence to be amplified is contained in a mixture of complicated long-chain nucleic acid strands. A large molar excess is preferred to improve the efficiency of the process.

The deoxyribonucleoside triphosphates dATP, dCTP, dGTP and TTP are also added to the synthesis mixture in adequate amounts and the resulting solution is heated to about 90-100°C for from about 1 to 10 minutes, preferably from 1 to 4 minutes. After this heating period the solution is allowed to cool to room temperature, which is preferable for the primer hybridization. To the cooled mixture is added an agent for polymerization, and the reaction is allowed to occur under conditions known in the art. This synthesis reaction may occur at from room temperature up to a temperature above which the inducing agent no longer functions efficiently. Thus, for example, if an *E. coli* DNA polymerase is used as agent for polymerization, the temperature is generally no greater than about 40°C. Most conveniently the reaction occurs at room temperature.

The agent for polymerization of the nucleotide triphosphates may be any compound or system which will function to accomplish the synthesis of primer extension products, including enzymes. Suitable enzymes for this purpose include, for example, *E. coli* DNA Polymerase I, Klenow fragment of *E. coli* DNA polymerase T4 DNA polymerase, other available DNA polymerases, reverse transcriptase, and other enzymes, including heat-stable enzymes, which will facilitate combination of the nucleotides in the proper manner to form the primer extension products which are complementary to each nucleic acid strand. Generally, the synthesis will be initiated at the 3' end of each primer and proceed in the 5' direction along the template strand, until synthesis terminates, producing molecules of different lengths. There may be agents, however, which initiate synthesis at the 5' end and in the other direction, using the same process as described above.

The newly synthesized strand and its complementary nucleic acid strand form double-stranded molecule which is used in the succeeding steps of the process. In the next step, the strands of the double-stranded molecule are separated using any of the procedures described above to provide single-stranded molecules.

New nucleic acid is synthesized on the single-stranded molecules. Additional agent for polymerization, nucleotides and primers may be added if necessary for the reaction to proceed under the conditions prescribed above. Again, the synthesis will be initiated at one end of the oligonucleotide primers and will proceed along the single strands of the template to produce additional nucleic acid. After this step, half of the extension product will consist of the specific nucleic acid sequence bounded by the two primers.

The steps of strand separation and extension product synthesis can be repeated as often as needed to produce the desired quantity of the specific nucleic acid sequence. As will be described in further detail below, the amount of the specific nucleic acid sequence produced will accumulate in an exponential fashion.

When it is desired to produce more than one specific nucleic acid sequence from the first nucleic acid or mixture of nucleic acids, the appropriate number of different oligonucleotide primers are utilized. For example, if two different specific nucleic acid sequences are to be produced, four primers are utilized. Two of the primers are specific for one of the specific nucleic acid sequences and the other two primers are specific for the second specific nucleic acid sequence. In this manner, each of the two different specific sequences can be produced exponentially by the present process.

The present invention can be performed in a step-wise fashion where after each step new reagents are added, or simultaneously, where all reagents are added at the initial step, or partially step-wise and partially simultaneous, where fresh reagent is added after a given number of steps. If a method of strand separation, such as heat, is employed which will inactivate the inducing agent, as in the case of a heat-labile enzyme then it is necessary to replenish the agent for polymerization after every strand separation step. The simultaneous method may be utilized when an enzymatic means is used for the strand separation step. In the simultaneous procedure, the reaction mixture may contain, in addition to the nucleic acid strand(s) containing the desired sequence, the strand-separating enzyme (e.g., helicase), an appropriate energy source for the rating enzyme, such as rATP, the four nucleotides, the oligonucleotide primers in molar excess, and the inducing agent, e.g., Klenow fragment of *E. coli* DNA Polymerase I.

If heat is used for denaturation in a simultaneous process, a heat-stable enzyme such as a thermostable polymerase may be employed which will operate at an elevated temperature, preferably 65-90°C depending on the agent for

polymerization, at which temperature the nucleic acid will consist of single and double strands in equilibrium. For smaller lengths of nucleic acid, lower temperatures of about 50°C may be employed. The upper temperature will depend on the temperature at which the enzyme will degrade or the temperature above which an insufficient level of primer hybridization will occur. Such a heat-stable enzyme is described, e.g., by A. S. Kaledin et al., *Biokhimiya*, 45 644-651 (1980). Each step of the process will occur sequentially notwithstanding the initial presence of all the reagents. Additional materials may be added as necessary. After the appropriate length of time has passed to produce the desired amount of the specific nucleic acid sequence, the reaction may be halted by inactivating the enzymes in any known manner or separating the components of the reaction.

In an alternative method using a thermostable enzyme, in step (a) the primers, enzyme and nucleotide triphosphates are contacted with the nucleic acid sample. In step (b) the mixture is treated to denature the nucleic acids and then incubated at a temperature at which the primers can hybridize to complementary sequences in the nucleic acid sample. In step (c) the mixture is heated for an effective time and at an effective temperature to promote the activity of the enzyme, and to synthesize, for each different sequence being amplified, an extension product of each primer which is complementary to each nucleic acid strand template, but not so high as to separate each extension product from its complementary strand template.

In step (d) the mixture is heated for an effective time and at an effective temperature to separate the primer extension products from the templates on which they were synthesized to produce single-stranded molecules, but not so high as to denature the enzyme irreversibly. In step (e) the mixture is cooled for an effective time and to an effective temperature to promote hybridization of each primer to each of the single-stranded molecules produced in step (d). Finally, in step (f) the mixture is heated for an effective time and to an effective temperature to promote the activity of the enzyme and to synthesize, for each different sequence being amplified, an extension product of each primer which is complementary to each nucleic acid strand template produced in step (d), but not so high as to separate each extension product from its complementary strand template, where steps (e) and (f) may be carried out simultaneously or sequentially.

A preferred thermostable enzyme which may be employed in the process herein is extracted and purified from *Thermus aquaticus* and has a molecular weight of about 86,000-90,000 daltons. This enzyme is more fully described in Example VIII hereinbelow.

The process of the present invention may be conducted continuously. In one preferred embodiment of an automated process wherein a thermostable enzyme is employed, the reaction may be cycled through a denaturing region, a primer annealing region, and a reaction region. In another embodiment, the enzyme used for the synthesis of primer extension products can be immobilized in a column. The other reaction components can be continuously circulated by a pump through the column and a heating coil in series, thus the nucleic acids produced can be repeatedly denatured without inactivating the enzyme.

In one preferred embodiment even where a thermostable enzyme is not employed and the temperature is raised and lowered, one such instrument is an automated machine for handling the amplification reaction of this invention. Briefly, this instrument utilizes a liquid handling system under computer control to make liquid transfers of enzyme stored at a controlled temperature in a first receptacle into a second receptacle whose temperature is controlled by the computer to conform to a certain incubation profile. The second receptacle stores the nucleic acid sequence(s) to be amplified plus the nucleotide triphosphates and primers. The computer includes a user interface through which a user can enter process parameters which control the characteristics of the various steps in the amplification sequence such as the times and temperatures of incubation, the amount of enzyme to transfer, etc.

A preferred machine which may be employed which is specifically adapted for use with a thermostable enzyme utilizes temperature cycling without a liquid handling system because the enzyme need not be transferred at every cycle. Briefly, this instrument consists of the following systems:

1. A heat-conducting container for holding a given number of tubes, preferably 500  $\mu$ l tubes, which contain the reaction mixture of nucleotide triphosphates, primers, nucleic acid sequences, and enzyme.
2. A means to heat, cool, and maintain the heat-conducting container above and below room temperature, which means has an input for receiving a control signal for controlling which of the temperatures at or to which the container is heated, cooled or maintained. (This may be Peltier heat pumps available from Materials Electronics Products Corporation in Trenton, N.J. or a water heat exchanger.)
3. A computer means (e.g., a microprocessor controller), coupled to the input of said means, to generate the signals which control automatically the amplification sequence, the temperature levels, and the temperature ramping and timing.

The present invention is demonstrated diagrammatically below where double-stranded DNA containing the desired sequence [S] comprised of complementary strands [S<sup>+</sup>] and [S<sup>-</sup>] is utilized as the nucleic acid. During the first and each subsequent reaction cycle extension of each oligonucleotide primer on the original template will produce one new

ssDNA molecule product of indefinite length which terminates with only one of the primers. These products, hereafter referred to as "long products," will accumulate in a linear fashion; that is, the amount present after any number of cycles will be proportional to the number of cycles.


The long products thus produced will act as templates for one or the other of the oligonucleotide primers during subsequent cycles and will produce molecules of the desired sequence [S<sup>+</sup>] or [S<sup>-</sup>]. These molecules will also function as templates for one or the other of the oligonucleotide primers, producing further [S<sup>+</sup>] and [S<sup>-</sup>], and thus a chain reaction can be sustained which will result in the accumulation of [S] at an exponential rate relative to the number of cycles.

By-products formed by oligonucleotide hybridizations other than those intended are not self-catalytic (except in rare instances) and thus accumulate at a linear rate.

The specific sequence to be amplified, [S], can be depicted diagrammatically as:




The appropriate oligonucleotide primers would be:



so that if DNA containing [S]




is separated into single strands and its single strands are hybridized to Primers 1 and 2, the following extension reactions can be catalyzed by DNA polymerase in the presence of the four deoxyribonucleoside triphosphates:




On denaturation of the two duplexes formed, the products are:





If these four strands are allowed to rehybridize with Primers 1 and 2 in the next cycle, the agent for polymerization will catalyze the following reactions:



If the strands of the above four duplexes are separated, the following strands are found:

55

5' AAAAAAAAAAXXXXXXXXXXCCCCCCCCC 3'  
newly synthesized [5']

3'....zzzzzzzzzzzzzzzzzzzzTTTTTTTTTYYYYYYYYYGGGGGGGGGG 5'  
first cycle synthesized long product 1

3'....zzzzzzzzzzzzzzzzzzzzTTTTTTTTTYYYYYYYYYGGGGGGGGGG 5'  
newly synthesized long product 1

5'...zzzzzzzzzzzzzzzzzzzzAAAAAAAAXXXXXXXXXXCCCCCCCCCzzzzzzzzz...3'  
original template strand

5' AAAAAAAAAAXXXXXXXXXXCCCCCCCCCzzzzzzzzzzzzzzzzzzzz...3'  
newly synthesized long product 2

3'..zzzzzzzzzzzzzzzzTTTTTTTTTTTTYYYYYYYYYYGGGGGGGGGGzzzzzzzzzzzzzzzz..5'  
original template strand

3' TTTT TTTT TTTT YYYYYYYYYY GGGGGGGGGG 5'  
newly synthesized [S<sup>-</sup>]

5' AAAAAAAAAAXXXXXXXXXXCCCCCCCCCzzzzzzzzzzzzzzzz...3'  
first cycle synthesized long product 2

It is seen that each strand which terminates with the oligonucleotide sequence of one primer and the complementary sequence of the other is the specific nucleic acid sequence [S] that is desired to be produced.

The steps of this process can be repeated indefinitely, being limited only by the amount of Primers 1 and 2, agent for polymerization and nucleotides present. The amount of original nucleic acid remains constant in the entire process, because it is not replicated. The amount of the long products increases linearly because they are produced only from the original nucleic acid. The amount of the specific sequence increases exponentially. Thus, the specific sequence will become the predominant species. This is illustrated in the following table, which indicates the relative amounts of the species theoretically present after  $n$  cycles, assuming 100% efficiency at each cycle:

Number of Double Strands After 0 to n Cycles			
Cycle Number	Template	Long Products	Specific Sequence [S]
0	1	-	-
1	1	1	0
2	1	2	1
3	1	3	4
5	1	5	26
10	1	10	1013
15	1	15	32,752
20	1	20	1,048,555
n	1	n	$(2^n - n - 1)$

When a single-stranded nucleic acid is utilized as the template, only one long product is formed per cycle.

The desired amount of cycles of this reaction will depend on, e.g., the nature of the sample. Fewer cycles will be required if the sample being analyzed is pure. If the sample is a complex mixture of nucleic acids, more cycles will be required to amplify the signal sufficiently for it to be detected by the method herein. For human genomic DNA preferably 15-30 cycles are carried out to amplify the sequence sufficiently that a clearly detectable signal is produced (i.e., so that background noise does not interfere with detection).

In one embodiment of the invention herein, the amplified sample suspected of containing the sequence variation, whether resulting from cancer, an infectious disease, a genetic disease, or just normal genetic polymorphism, is spotted directly on a series of membranes and each membrane is hybridized with a different labeled sequence-specific oligo-

nucleotide probe. One procedure for spotting the sample on a membrane is described by Kafotos et al., Nucleic Acids Research, 7:1541-1552 (1979).

Briefly, the DNA sample affixed to the membrane may be pretreated with a prehybridization solution containing sodium dodecyl sulfate, Ficoll, serum albumin and various salts prior to the probe being added. Then, a labeled oligonucleotide probe which is specific to each sequence to be detected is added to a hybridization solution similar to the prehybridization solution. The hybridization solution is applied to the membrane and the membrane is subjected to hybridization conditions that will depend on the probe type and length, type and concentration of ingredients, etc. Generally, hybridization is carried out at about 25-75°C, preferably 35 to 85°C, for 0.25-50 hours, preferably less than three hours. The greater the stringency of conditions, the greater the required complementarity for hybridization between the probe and sample. If the background level is high, stringency may be increased accordingly. The stringency can also be incorporated in the wash.

After the hybridization the sample is washed of unhybridized probe using any suitable means such as by washing one or more times with varying concentrations of standard saline phosphate EDTA (SSPE) (180 mM NaCl, 10 mM NaCl, 10 mM NaHPO<sub>4</sub> and 1 M EDTA, pH 7.4) solutions at 25-75°C for about 10 minutes to one hour, depending on the temperature. The label is then detected by using any appropriate detection techniques.

The sequence-specific oligonucleotide employed herein is an oligonucleotide which is generally prepared and selected as described above for preparing and selecting the primers. The sequence-specific oligonucleotide must encompass the region of the sequence which spans the nucleotide variation being detected and must be specific for the nucleotide variation being detected. For example, if it is desired to detect whether a sample contains the mutation for sickle cell anemia, one oligonucleotide will be prepared which contains the nucleotide sequence site characteristic of the normal  $\beta$ -globin gene, and one oligonucleotide will be prepared which contains the nucleotide sequence characteristic of the sickle cell allele. Each oligonucleotide would be hybridized to duplicates of the same sample to determine whether the sample contains the mutation.

The polymorphic areas of HLA class II genes are localized to specific regions of the second exon and are flanked by conserved sequences, so that oligonucleotide primers complementary to opposite strands of the conserved 5' and 3' ends of the second exon can be prepared.

The number of oligonucleotides employed for detection of the polymorphic areas of the HLA class II genes will vary depending on the type of gene, which has regions of base pair variation which may be clustered or spread apart. If the regions are clustered, as in the case with HLA-DQ $\alpha$ , then one oligonucleotide is employed for each allele. If the regions are spread apart, as in the case with HLA-DQ $\beta$  and HLA-DR $\beta$ , then more than one probe, each encompassing an allelic variant, will be used for each allele. In the case of HLA-DQ $\beta$  and HLA-DR $\beta$ , three probes are employed for the three regions of the locus where allelic variations may occur. For detection of sequences associated with insulin-dependent diabetes mellitus (IDDM) four probes for the HLA-DR $\beta$  second exon are employed.

Haplotypes can be inferred from segregation analysis in families or, in some cases, by direct analysis of the individual DNA sample. Specific allelic combinations (haplotypes) of sequence-specific oligonucleotide reactivities can be identified in heterozygous cells by using restriction enzyme digestion of the genomic DNA prior to amplification.

For example, if in DQ $\beta$  one finds three highly variable subregions A, B, and C within a single amplified region, and if there are six different sequences at each region (A1-6, B1-6, C1-6), then an individual could be typed in the DQ $\beta$  locus by sequence-specific oligonucleotide probe analysis as containing A1, A2; B2 B3; C1, C4, with the possible haplotype combinations of A1, B2, C1; A1, B2, C4; A2, B2, C1; A2, B2, C4; A1, B3, C1; A1, B3, C4; A1, B2, C1; and A1, B2, C4.

If the genomic DNA is digested with a polymorphic restriction enzyme prior to amplification, and if the enzyme cuts both alleles between the primers, there is no reactivity with the sequence-specific probes due to lack of amplification, and the result is uninformative. If the enzyme cuts neither allele, the probe results with digested and undigested genomic DNA are the same and the result is uninformative. If the enzyme cuts only one allele, however, then one can infer both haplotypes by comparing the probe reactivity patterns on digested and undigested DNA.

The haplotypes can be deduced by comparing sequence-specific oligonucleotide reactivities with uncut genomic DNA and genomic DNA cut with one or several enzymes known to be polymorphic and to recognize sites between the primers.

The length of the sequence-specific oligonucleotide will depend on many factors, including the particular target molecule being detected, the source of oligonucleotide, and the nucleotide composition. For purposes herein, the probe typically contains 15-25 nucleotides, although it may contain more or fewer nucleotides. While oligonucleotides which are at least 19-mers in length may enhance specificity and/or sensitivity, probes which are less than 19-mers, e.g., 16-mers, show more sequence-specific discrimination, presumably because a single mismatch is more destabilizing. Because amplification increases specificity so that a longer length is less critical, and hybridization and washing temperatures can be lowered for the same salt concentration, it is preferred to use probes which are less than 19-mers.

Where the sample is first placed on the membrane and then detected with the oligonucleotide, the oligonucleotide must be labeled with a suitable label moiety, which may be detected by spectroscopic, photochemical, biochemical,

immunochemical or chemical means. Immunochemical means include antibodies which are capable of forming a complex with the oligonucleotide under suitable conditions, and biochemical means include polypeptides or lectins capable of forming a complex with the oligonucleotide under the appropriate conditions. Examples include fluorescent dyes, electron-dense reagents, enzymes capable of depositing insoluble reaction products or being detected chronogenically, such as alkaline phosphatase, a radioactive label such as  $^{32}\text{P}$ , or biotin. If biotin is employed, a spacer arm may be utilized to attach it to the oligonucleotide.

Alternatively, in one "reverse" dot blot format, at least one of the primers and/or at least one of the four nucleotide triphosphates is labeled with a detectable label, so that the resulting amplified sequence is labeled. These labeled moieties may be present initially in the reaction mixture or added during a later cycle. Then an unlabeled sequence-specific oligonucleotide capable of hybridizing with the amplified nucleic acid sequence, if the variation(s) in sequence (whether normal or mutant) is/are present, is spotted on (affixed to) the membrane under prehybridization conditions as described above. The amplified sample is then added to the pretreated membrane under hybridization conditions as described above. Finally, detection means are used to determine if an amplified sequence in the nucleic acid sample has hybridized to the oligonucleotide affixed to the membrane. Hybridization will occur only if the membrane-bound sequence containing the variation is present in the amplification product, i.e. only if the sequence of the probe is complementary to a region of the amplified sequence.

In another version of the "reverse" dot blot format, the amplification is carried out without employing a label as with the "forward" dot blot format described above, and a labeled sequence-specific oligonucleotide probe capable of hybridizing with the amplified nucleic acid sequence only if the sequence of the probe is complementary to a region of the amplified nucleic acid is spotted on (affixed to) the membrane under prehybridization conditions as described above. The amplified sample is then added to the pretreated membrane under hybridization conditions as described above. Then the labeled oligonucleotide or a fragment thereof is released from the membrane in such a way that a detection means can be used to determine if an amplified sequence in the sample hybridized to the labeled oligonucleotide. The release may take place, for example, by adding a restriction enzyme to the membrane which recognizes a restriction site in the probe. This procedure, known as oligomer restriction, is described more fully in EP Patent Publication 164,054 published December 11, 1985.

For purposes of this invention, the genetic diseases which may be detected include specific deletions, insertions and/or substitutions in any base pair mutation or polymorphism in nucleic acids, for example, genomic DNA, from any organism. Examples of diseases in which a base pair variation is known include sickle cell anemia, hemoglobin C disease,  $\alpha$ -thalassemia,  $\beta$ -thalassemia, and the like. Other diseases that may be detected include cancerous diseases such as those involving the RAS oncogenes, e.g., the n-RAS oncogene, and infectious diseases.

The process herein may also be used for HLA typing in the areas of tissue transplantation, disease susceptibility, and paternity determination. The HLA class II genes, consisting of the  $\alpha$  and  $\beta$  genes from the HLA-DR, HLA-DQ and HLA-DP regions, are highly polymorphic; their genetic complexity at the DNA level is significantly greater than the polymorphism currently defined by serological typing. In addition, the process herein may be employed to detect certain DNA sequences coding for HLA class II  $\beta$  proteins (e.g., DR $\beta$ ) associated with insulin-dependent diabetes mellitus (IDDM). Briefly, the four DNA sequences associated with IDDM are selected from the group consisting of:

- 1) 5'-GAGCTGCGTAAGTCTGAG-3',
- 2) 5'-GAGGAGTTCTCGCTTC-3',
- 3) 5'-CCTGTCGCCGAGTCCTGG-3', and
- 4) 5'-GACATCCTGGAAGACGAGAGA-3',

or the DNA strands that are complementary thereto. Sequence-specific probes may be prepared that will hybridize to one or more of these sequences.

The invention herein also contemplates a kit format which comprises a packaged multicontainer unit having containers for each primer and probe utilized, a container with the agent for polymerization to synthesize the primer extension products, such as enzymes, a container for each of the four nucleotides, and a container with means to detect the label (such as an avidin-enzyme conjugate and enzyme substrate and chromogen if the label is biotin). In addition, the kit may have a container which includes a positive control for the probe containing one or more nucleic acids with the nucleotide variation to be detected and a negative control for the probe which does not contain the nucleic acids with the nucleotide variation to be detected. Moreover, the kit may also have a container with means for separating the strands of any double-stranded nucleic acids contained in the sample such as a helicase or sodium hydroxide.

The following examples illustrate various embodiments of the invention and are not intended to be limiting in any respect. In the examples all parts and percentages are by weight if solid and by volume if liquid, and all temperatures are in degrees Celsius, unless otherwise indicated.

**EXAMPLE I**

This example illustrates how the process herein can be used to distinguish normal alleles (A) from sickle cell alleles (S) from Hemoglobin C disease alleles (C).

**1. Synthesis of the Primers**

The following two oligonucleotide primers were prepared by the method described below:



These primers, both 20-mers, anneal to opposite strands of the genomic DNA with their 5' ends separated by a distance of 110 base pairs.

A. Automated Synthesis Procedures: The diethylphosphoramidites, synthesized according to Beaucage and Caruthers (Tetrahedron Letters (1981) 22:1859-1862) were sequentially condensed to a nucleoside derivatized controlled pore glass support. The procedure included detritylation with trichloroacetic acid in dichloromethane, condensation using benzotriazole as activating proton donor, and capping with acetic anhydride and dimethylaminopyridine in tetrahydrofuran and pyridine. Cycle time was approximately 30 minutes. Yields at each step were essentially quantitative and were determined by collection and spectroscopic examination of the dimethoxytrityl alcohol released during detritylation.

B. Oligodeoxyribonucleotide Deprotection and Purification Procedures: The solid support was removed from the column and exposed to 1 ml concentrated ammonium hydroxide at room temperature for four hours in a closed tube. The support was then removed by filtration and the solution containing the partially protected oligodeoxynucleotide was brought to 55°C for five hours. Ammonia was removed and the residue was applied to a preparative polyacrylamide gel. Electrophoresis was carried out at 30 volts/cm for 90 minutes after which the band containing the product was identified by UV shadowing of a fluorescent plate. The band was excised and eluted with 1 ml distilled water overnight at 4°C. This solution was applied to a column and eluted with a 7-13% gradient of acetonitrile in 1% ammonium acetate buffer at pH 6.0. The elution was monitored by UV absorbance at 260 nm and the appropriate fraction collected, quantitated by UV absorbance in a fixed volume and evaporated to dryness at room temperature in a vacuum centrifuge.

C. Characterization of Oligodeoxyribonucleotides: Test aliquots of the purified oligonucleotides were  $^{32}\text{P}$  labeled with polynucleotide kinase and  $\gamma\text{-}^{32}\text{P}\text{-ATP}$ . The labeled compounds were examined by autoradiography of 14-20% polyacryl amide gels after electrophoresis for 45 minutes at 50 volts/cm. This procedure verifies the molecular weight. Base composition was determined by digestion of the oligodeoxyribonucleotide to nucleosides by use of venom diesterase and bacterial alkaline phosphatase and subsequent separation and quantitation of the derived nucleosides using a reverse phase HPLC column and a 10% acetonitrile, 1% ammonium acetate mobile phase.

**II. Isolation of Human Genomic DNA from Cell Line**

High molecular weight genomic DNA was isolated from the lymphoid cell line GM2064 using essentially the method of Maniatis et al., Molecular Cloning (1982), 280-281. GM2064 ( Human Mutant Cell Repository, Camden, N.J.) was originally isolated from an individual homozygous for hereditary persistence of fetal hemoglobin (HPFH) and contains no  $\beta$ - or  $\delta$ -globin gene sequences. This cell line was maintained in RPMI-1640 with 10% fetal calf serum.

**III. Isolation of Human Genomic DNA from Clinical Samples**

Five clinical blood samples designated AA (from a known normal individual), AS (from a known sickle cell carrier), SS (from a known sickle cell individual), SC (from a known sickle cell/hemoglobin C diseased individual), and AC (from a known hemoglobin C disease carrier) were obtained from Dr. Bertram Lubin of Children's Hospital in Oakland, California. One clinical DNA sample designated CC (from a known hemoglobin C diseased individual) was obtained from Dr. Stephen Embury of San Francisco General Hospital in San Francisco, California.

Genomic DNA from the first five of these samples was prepared from the buffy coat fraction, which is composed primarily of peripheral blood lymphocytes, as described by Saiki et al., Biotechnology, 3: 1008-1012 (1985).

## IV. Amplification Reaction

One microgram of DNA from each of the seven DNA samples (10  $\mu$ l of 100  $\mu$ g/ml DNA) was amplified in an initial 100  $\mu$ l reaction volume containing 10  $\mu$ l of a solution containing 100 mM Tris-HCl buffer (pH 7.5), 500 mM NaCl, and 100 mM  $MgCl_2$ , 10  $\mu$ l of 10  $\mu$ M of primer PCO3, 10  $\mu$ l of 10  $\mu$ M of primer PCO4, 15  $\mu$ l of 40 mM dNTP (contains 10 mM each of dATP, dCTP, dGTP and TTP), and 45  $\mu$ l of water.

Each reaction mixture was held in a heat block set at 95°C for 10 minutes to denature the DNA. Then each DNA sample underwent 25 cycles of amplification where each cycle was composed of four steps:

- (1) spin briefly (10-20 seconds) in microcentrifuge to pellet condensation and transfer the denatured material immediately to a heat block set at 30°C for two minutes to allow primers and genomic DNA to anneal,
- (2) add 2  $\mu$ l of a solution prepared by mixing 39  $\mu$ l of the Klenow fragment of *E. coli* DNA Polymerase I (New England Biolabs, 5 units/ $\mu$ l), 39  $\mu$ l of a salt mixture of 100 mM Tris buffer (pH 7.5), 500 mM NaCl and 100 mM  $MgCl_2$  and 312  $\mu$ l of water,
- (3) allowing the reaction to proceed for two minutes at 30°C, and
- (4) transferring the samples to the 95°C heat block for two minutes to denature the newly synthesized DNA, except this reaction was not carried out at the last cycle.


The final reaction volume was 150  $\mu$ l, and the reaction mixture was stored at -20°C.

## V. Synthesis and Phosphorylation of Oligodeoxyribonucleotide Probes

Three labeled DNA probes, designated RS17, RS18 and RS21, of the following sequences were prepared as follows:

5' - \*CTCCTAAGGAGAAGTCTGC - 3' (RS17)  
 5' - \*CTCCTGAGGAGAAGTCTGC - 3' (RS18)  
 5' - \*CTCCTGTGGAGAAGTCTGC - 3' (RS21)

where \* indicates the label. These probes are 19 bases long and span the fifth through eleventh codons of the gene. RS18 is complementary to the normal  $\beta$ -globin allele ( $\beta^A$ ), RS21 to the sickle cell anemia allele ( $\beta^S$ ), and RS17 to the hemoglobin C disease allele ( $\beta^C$ ). RS17 and RS21 differ from RS18 by a single base. The schematic diagram of primers and probes is given below:



These three probes were synthesized according to the procedures described in Section I. The probes were labeled by contacting 10 pmole thereof with 4 units of T4 polynucleotide kinase (New England Biolabs) and about 40 pmole  $\gamma$ -<sup>32</sup>P-ATP (New England Nuclear, about 7000 Ci/mmol) in a 40  $\mu$ l reaction volume containing 70 mM Tris buffer (pH 7.6), 10 mM  $MgCl_2$ , 1.5 mM spermine, 100 mM dithiothreitol and water for 60 minutes at 37°C. The total volume was then adjusted to 100  $\mu$ l with 25 mM EDTA and purified according to the procedure of Maniatis et al., *Molecular Cloning* (1982), 466-467 over a 1 ml Bio Gel P-4 (BioRad) spin dialysis column equilibrated with Tris-EDTA (TE) buffer (10 mM Tris buffer, 0.1 mM EDTA, pH 8.0). TCA precipitation of the reaction product indicated that for RS17 the specific activity was 5.2  $\mu$ Ci/pmol and the final concentration was 0.118 pmol/ $\mu$ l. For RS18 the specific activity was 4.6  $\mu$ Ci/pmol and the final concentration was 0.114 pmol/ $\mu$ l. For RS21 the specific activity was 3.8  $\mu$ Ci/pmol and the final concentration was 0.112 pmol/ $\mu$ l.

## VI. Dot Blot Hybridizations

Five microliters of each of the 150  $\mu$ l amplified samples from Section III was diluted with 195  $\mu$ l 0.4 N NaOH, 25

mM EDTA and spotted onto three replicate cationic nylon filters by first wetting the filter with water, placing it in an apparatus for preparing dot blots which holds the filter in place, applying the samples, and rinsing each well with 0.4 ml of 20 x SSPE (3.6 M NaCl 200 mM NaH<sub>2</sub>PO<sub>4</sub>, 20 mM EDTA), as disclosed by Reed and Mann, Nucleic Acids Research, 13, 7202-7221 (1985). The filters were then removed, rinsed in 20 x SSPE, and baked for 30 minutes at 80°C in a vacuum oven.

After baking, each filter was then contacted with 6 ml of a hybridization solution consisting of 5 x SSPE, 5 x Denhardt's solution (1 x = 0.02% polyvinylpyrrolidone, 0.02% Ficoll, 0.02% bovine serum albumin, 0.2 mM Tris-HCl, 0.2 mM EDTA, pH 8.0) and 0.5% SDS and incubated for 60 minutes at 55°C. Then 5 µl each of probes RS17, RS18 and RS21 was added to the hybridization solution and the filter was incubated for 60 minutes at 55°C.

Finally, each hybridized filter was washed twice with 100 ml 2 x SSPE and 0.1% SDS for 10 minutes at room temperature. As a third wash for RS17, 250 ml of 5 x SSPE and 0.1% SDS was added and the filter heated for five minutes at 55°C. For RS18 and 21, the filters were treated with 250 ml of 4 x SSPE, 0.1% SDS for five minutes at 55°C. There was a faint background with RS18 and RS21 because 4 x SSPE at 55°C was not sufficiently stringent. The washing with the RS18 and RS21 probes was repeated in 250 ml of 5 x SSPE, 0.1% SDS for three minutes at 55°C. There was no change in the background. The wash was repeated with 5 x SSPE at 60°C for one minute and an additional wash of the same stringency was done for three minutes. This resulted in virtually no background. The genotypes were readily apparent after 90 minutes of autoradiography.

## VII. Discussion of Autoradiogram

The autoradiogram of the dot blot of the seven amplified genomic DNA samples hybridized with allele-specific β-globin probes RS18, RS21 and RS17 was analyzed after 12 hours. The negative control GM2064 was included. The results clearly indicate that each allele-specific probe annealed only to the DNA samples which had at least one copy of the β-globin allele to which it was perfectly matched. For example, the β<sup>A</sup>-specific probe, RS18, hybridized only to samples AA (β<sup>A</sup>β<sup>A</sup>), AS'(β<sup>A</sup>β<sup>C</sup>), and AC (β<sup>A</sup>β<sup>C</sup>).

### EXAMPLE II

To determine the minimum levels of detection by dot blot, eight serial dilutions containing 128, 64, 32, 16, 8, 4, 2 and 1 ng of normal genomic DNA were made from sample AA and subjected to 25 cycles of amplification as described in Example I.

As controls, the amplified samples of AA and SS from Example I were similarly diluted as well. A total of 75 µl (one-half) of each sample was mixed with 125 µl of 0.65 N NaOH and 25 mM EDTA and the mixture was applied to a nylon filter. Then the filter was rinsed in 20 x SSPE and baked for 30 minutes at 80°C.

The filter was then probed as described in Example I with RS18 (the β<sup>A</sup> probe) to determine the detection threshold. The prehybridization solution was 8 ml of 5 x SSPE, 5 x Denhardt's solution, 0.5% SDS for 40 minutes at 55°C and the hybridization solution was the same plus 10 µl of RS18 for 80 minutes at 55°C. The filters were then washed with 2 x SSPE, 0.1% SDS for 10 minutes at room temperature twice and then with 5 x SSPE, 0.1% SDS for three minutes at 60°C.

The autoradiogram obtained after hybridization with RS18 after 17 hours of exposure revealed positive signals in all samples containing the AA DNA. The SS sample was visible after 17 hours but the intensity of the 64 ng SS was equivalent to the intensity of 1 ng AA, which is a signal-to-noise ratio of 64:1. The intensity of the signal present in the 0.5 ng spot suggested that amplification of samples containing significantly less than 1 ng is possible. (One nanogram is the amount of genomic DNA present in 150 diploid cells.)

### EXAMPLE III

#### A. Amplification and Detection of HLA-DQα Sequences

##### I. Preparation of Primers

Oligonucleotides designated GH26 and GH27 complementary to opposite strands of the conserved 5' and 3' ends of the DQα second exon were used as primers to amplify a 240 base pair fragment. The primers, having the following sequences, were prepared as described in Example I.

5'-GTGCTGCAGGTGTAACTTGTACCAG-3' (GH26)  
 5'-CACGGATCCGGTAGCAGCGGTAGATTG-3' (GH27)

## II. Preparation of Probes

Based on the analysis of HLA-DQ $\alpha$  sequences from diverse sources, which were grouped into allelic variants, the following probes from variable regions of the DQ $\alpha$  second exon encompassing each variant were synthesized and labeled as described in Example I. The two variable regions of the HLA-DQ $\alpha$  second exon (called "exon I"), segments A and B, are shown in Table I. The entire scheme of primers (designated by PCR $\rightarrow$ ), probes and HLA-DQ $\alpha$  sequence is shown in Table II, and the amino acid abbreviations used therein are shown in Table III.

TABLE I

### HLA DQ $\alpha$ (segment A):

	35	40	
	GlyAspGluGluPheTyrValAspLeuGluArgLysGlu		
	GGAGATGAGGAGTTCTACGTGGACCTGGAGAGGAAGGAG		DR1,2,w6
	-----C-----A-----		DR5,8
	-----C-----T-----		DR4,7,9
	-----C-----G-----		DR3
25	DXA: -----C-----T-----C-----A-----		

### HLA DQ $\alpha$ (segment B):

	45	50	55	60	
	AlaTrpArgTrpProGluPheSerLysPheGlyGlyPheAspProGlnGly				
	GCCTGGCGGTGGCCTGAGTTCAGCAAATTTGGAGGTTTTGACCCGCAGGGT				DR1,2,5,w6,8
	-T-----A-T-----CT-----C-----G-----A-A-A-----ATT-				DR4,9
	-T-----AA-T-----CT-----CA-G-C-----A-----A-----ATT-				DR7
	-T-----T-T-T-----TTC-----AC-----A-----A-----ATT-				DR3
35	DXA: -T-----A--T-----AT-----T-----AT-A-----A--				



**Exon-1:**

PCR →      ← PCR

### Amino Acid Abbreviation Codes

Amino Acid Abbreviation Codes		
Alanine	Ala	A
Arginine	Arg	R
Asparagine	Asn	N

TABLE III (continued)

Amino Acid Abbreviation Codes		
Aspartic Acid	Asp	D
Cysteine	Cys	C
Glutamine	Gln	Q
Glutamic Acid	Glu	E
Glycine	Gly	G
Histidine	His	H
Isoleucine	Ile	I
Leucine	Leu	L
Lysine	Lys	K
Methionine	Met	M
Phenylalanine	Phe	F
Proline	Pro	P
Serine	Ser	S
Threonine	Thr	T
Tryptophan	Trp	W
Tyrosine	Tyr	Y
Valine	Val	V

The probes are as follows, where \* is the label.

5' -\*TGTTCCTGTTCTCAGAC-3' (GH66)  
 5' -\*TTCCGCAGATTAGAAAGAT-3' (GH67)  
 5' -\*TTCCACAGACTTAGATTTG-3' (GH68)  
 5' -\*CTCAGGCCACGCCAGGCA-3' (GH75)

The GH75 probe was derived from the DRw6 sequence, which is described by Auffray et al., *Nature*, 308:327 (1984). The GH67 probe was derived from the DR4 sequence, which is described by Auffray et al., *PNAS*, 79:6337 (1982). The GH68 probe was derived from the DR7 sequence which is described by Chang et al., *Nature*, 305:813 (1983). The GH66 probe was derived from the DR3 sequence, which is described by Schenning et al., *EMBO.J.*, 3: 447 (1984). A control oligonucleotide was derived from a conserved segment of all of these alleles.

### III. Origin and Preparation of Genomic DNA

Eleven DNA samples described below were prepared for subsequent amplification as described in Example I.

LG2 cell line (genotype DR1) from Drs. John Bell and Dan Denny of Stanford University, Stanford, California  
 PGF cell line (genotype DR2) from Drs. John Bell and Dan Denny  
 AVL cell line (genotype DR3) from Drs. John Bell and Dan Denny  
 DKB cell line (genotype DR4) from Drs. John Bell and Dan tinny  
 JGL cell line (genotype DR5) from Dr. Gerry Nepom of Virginia Mason Hospital, Seattle, Washington  
 APD cell line (genotype DR6) from Drs. John Bell and Dan Denny  
 LBF cell line (genotype DR7) from Drs. John Bell and Dan tinny  
 TAB cell line (genotype DR8) from Dr. Gerry Nepom  
 KOZ cell line (genotype DR9) from Dr. Gerry Nepom Sample GM2741A (genotype DR1,3) available from the Human Genetic Mutant Cell Repository, Camden, N.J.  
 Sample GM2676 (genotype DR4,3) available from the Human Genetic Mutant Cell Repository

### IV. Amplification Reaction

Each genomic DNA sample was amplified as described in Example I using GH26 and GH27 as primers for 28 cycles, except that the polymerization step 3 was carried out at 37°C in the presence of 10% by weight dimethylsulfoxide (DMSO), and except that the amplification reaction took place in the aluminum heating block of the above-described

automated liquid handling temperature-cycling instrument using the following program:

- 1) 2.5 min., 37°C to 98°C ramp (denature);
- 2) 3.0 min., 98°C to 37°C ramp (anneal);
- 3) add 1 unit Klenow fragment; and
- 4) 2.0 min., 37°C maintain (extend).

#### V. Dot Blot Hybridization

For each DNA sample, four duplicate nylon filters were spotted with 5 µl of the 150 µl amplified genomic DNA and one of probes GH66, 67, 68 or 75 was applied thereto as described in Example I except that DNA samples were neutralized before application to a nylon filter membrane, using a prehybridization solution of 6 x SSPE, 10 x Denhardt's solution and 0.5% SDS for one hour at 50°C and the same solution overnight at 50°C. The filters were washed with 0.1 x SSPE, 0.1% SDS for 10-15 minutes at 37°C. The filters were treated as described in Example I to obtain an autoradiogram.

#### VI. Discussion of Autoradiogram

The autoradiogram of the dot blot shows that the four HLA-DQα allele-specific probes, GH66, GH67, GH68, and GH75 complementary to four HLA-DQα allelic variants, may be used to define nucleotide sequence polymorphisms on amplified DNA from both homozygous and heterozygous individuals. The pattern of reactivity of the probe GH75 corresponds to the serologically defined type DQW1.

#### B. Amplification and Detection of HLA-DQβ Sequences

##### I. Preparation of Primers

Oligonucleotides designated GH28 and GH29 complementary to opposite strands of the conserved 5' and 3' ends of the DQ-β second exon were used as primers. The primers, having the following sequences, were prepared as described in Example I.

5' -CTCGGATCCGCATGTGCTACTTCACCAACG-3' (GH28)  
5' -GAGCTGCAGGTAGTTGTGTCTGCACAC-3' (GH29)

##### II. Preparation of Probes

Based on the analysis of HLA-DQβ sequences from diverse sources, which were grouped into allelic variants, the following probes from two variable regions of the DQβ second exon encompassing each variant were synthesized and labeled as described in Example I. The two regions, segments A (GH69-71) and B (GH60-62), as well as a third variable region, are shown in Table IV. The entire scheme of primers, probes and HLA-DQβ sequence is shown in Table V, where the amino acid abbreviations are shown in Table III above.

TABLE IV

HLA-DQ $\beta$  (segment A):

	20	25	30	
	GlyThrGluArgValArgGlyValThrArgHisIleTyr			
	GGGACGGAGCGCGTGC	GGGGTGTGACCAGACACATCTAT		DR1
10	-----TCT-----	-----T-----		DR2,4
	-----TCT-----	-----A-----		DR6
	-----C-----	-----T-----		DR8
	-----C-----	-----T-----		DR4'
	-----A-----	-----TCT-----	-----G-----	AG-----
15	DXB: -----	-----A-----	-----C-----	-----G-----
				-----T-----
				-----

HLA-DQ $\beta$  (segment B):

	45	50	55	60	
20	ValGlyValTyrArgAlaValThrProGlnGlyArgProValAlaGluTyrTrpAsn				
	GTGGGGGTGTACCGGGCAGTGACGCCGCGAGGGCGGCCTGTTGCCGAGTACTGGAAC				DR1
	-----C-----	-----G-----	-----A-----		DR2
	-----	-----G-----			DR6
25	-----T-----	-----S-----	-----T-----	-----C-----	CC-----
	-----A-----	-----G-----	-----T-----	-----C-----	AC-----
	-----T-----	-----G-----	-----T-----	-----T-----	AC-----
	-----A-----	-----T-----	-----G-----	-----T-----	CC-----
30	DXB: -----	-----T-----	-----A-----	-----T-----	-----A-----
					-----G-----
					-----CGA-----
					-----T-----
					-----AGCA-----
					-----C-----
					-----AG-----
					-----C-----
					-----

HLA-DQ $\beta$  (segment C):

	65	70	75	
35	GluValLeuGluGlyAlaArgAlaSerValAspArgVal			
	GAAGTCCTGGAGGGGGCCCGGGCGTCGGTGGACAGGGTG			DR1
	-----A-----	-----GA-----	-----T-----	-----C-----
	-----A-----	-----A-----	-----GA-----	-----T-----
	-----CA-----	-----A-----	-----A-----	-----CC-----
40	-----CA-----	-----A-----	-----AAA-----	-----G-----
	-----CT-----	-----T-----	-----CA-----	-----AG-----
				-----CG-----
				-----A-----
				-----

TABLE V  
Alignment of HLA-DQB Protein Sequence

Exon-1:	20	40	60	80																		
DC8PG:	DFVYQFKGMCYF	TNGTERVRL	VTRYIYNREEVARFSDVG	VYRAVTP	LGPPAAEYNNMSQKEVLER	TRAELDTVCRHNYQLE	RTTLQRR	(DR4)														
JoanP:								(DR4, 3)														
MMCC4:								(DR4)														
NIN:		Y		E				(DR4')														
ARC:		G			RLD		DI	ED-SV	(DR8)													
D096:		G		E		RLD	DI	ED-SV	(DR8, 7)													
DQBS3:		L	G			RLD	DI	ED-SV	(DR4, 1)													
DCBP1:		S	S	VV		EF	L	L	DI	K	AV-R	(DR3, 6)										
D096:		S	S	IV		EF	L	L	DI	K	AV-R	(DR7, 8)										
CMCC3:		S	S	IV		EF	L	L	DI	K	AV-R	(DR3)										
DCBS1:		S	S	V		EF	L	L	DI	K	AV-R	(DR3)										
DQBC:		L	G	H	V		Q	R	V		GA	SV-R	EVAY-GI	(DR1)								
DCBP2:		L	G	H	V		Q	R	V		GA	SV-R	EVAY-GI	(DR1, 6)								
TAB:			Y		DV		Q	R	D		L	DI		(DR8)								
DQBH:							Q	R	D		G		EVAF-GI	(DR2)								
PGF:							Q	R	D		G			(DR2)								
CMCC6:			H				Q	R	V				EVGY-GI	(DR6)								
DX8:		L	V		G	A				EFQ	E	RSI	D	NY	DF	QE	AV	K	EA	Q		
	PCR	→				A						B						C			←	PCR



The probes are as follows, where \* is the label and \*\* indicates the best probes.

5' - \*CGGCAGGCGGCCCCAGCGG-3' (GH60)\*\*  
 5' - \*CGGCAGGCGGCCCCAGCAG-3' (GH61)\*\*  
 5' - \*CAACAGGCGGCCCCCTGCGG-3' (GH62)  
 5' - \*GATGTGTCTGGTCACACCCCG-3' (GH69)\*\*  
 5' - \*GATGCTTCTGCTCACAAGACG-3' (GH70)  
 5' - \*GATGTATCTGGTCACAAGACG-3' (GH71)

## 10 II. Amplification and Dot Blot Hybridization

Using the method generally described in Example IIIA, the probes were found to have reasonable specificity for the portions of the allele being detected in genomic DNA samples.

## 15 C. Amplification and Detection of HLA-DR $\beta$ Sequences

### I. Preparation of Primers

20 Oligonucleotides designated GH46 and GH50 complementary to opposite strands of the conserved 5' and 3' ends of the DR $\beta$  second exon were used as primers. The primers, having the following sequences, were prepared as described in Example I.

5' -CCGGATCCTTCGTGTCCCCACAGCACG-3' (GH46)  
 5' -CTCCCAACCCCGTAGTTGTGTCTGCA-3' (GH50)

### II. Preparation of Probes

30 Based on the analysis of HLA-DR $\beta$  sequences from diverse sources, which were grouped into allelic variants, the following probes from two variable regions of the DR $\beta$  second exon encompassing each variant were synthesized and labeled as described in Example I. The two regions, segments A (GH56-59) and B (GH51), are shown in Table VI.

35

40

45

50

55

Table VI

HLA-DR $\beta$  (segment A)

	6	10	15	20	25	30	35	40	45
ArgPheLeuGluGlnValLysHisGluCysHisPhePheAsnGlyThrGluArg									
MM#1:	R	F	L	E	Q	V	K	H	C
DRBS2:	-	-	-	-	-	-	-	-	-
CM#15:	-	-	-	-	-	-	-	-	-
CM#4:	-	-	-	-	-	-	-	-	-
CM#6:	-	-	-	-	-	-	-	-	-
DRBC:	-	-	-	-	-	-	-	-	-
LG2#2:	-	-	-	-	-	-	-	-	-
DRBP4:	-	-	-	-	-	-	-	-	-

HLA-DR $\beta$  (segment B)

	25	30	35	40	45
ValArgPheLeuAspArgTyrPheTyrHisGlnGluGluTyrValArgPheAspSerAspVal					
MM#1:	V	R	F	L	D
DRBS2:	-	W	N	-	I
CM#15:	-	-	-	-	-
CM#4:	-	-	-	-	-
CM#5:	-	-	-	-	-
CM#6:	-	-	-	-	-
DRBC:	-	-	-	-	-
LG2#2:	-	Q	Y	-	N
DRBP4:	-	-	-	-	-

The probes are as follows, where \* is the label.

5'-\*CTGATCAGGTTCCACACTCG-3' (GH51)  
 5'-\*CAGACGTAGAGTACTCC-3' (GH56)  
 5'-\*CAGACTTACGCAGCTCC-3' (GH57)  
 5'-\*CAGACTTAAGCAGCTCC-3' (GH58)  
 5'-\*CATGTTTAACCTGCTCC-3' (GH59)

### III. Amplification and Dot Blot Hybridization

Using the method generally described in Example IIIA, the probes were found to have reasonable specificity for the portions of the allele being detected in genomic DNA samples.

### IV. Analysis of HLA-DR $\beta$ Sequences Associated With IDDM

Several HLA class II beta genes were isolated from clinical blood samples of diverse HLA-typed IDDM individuals (from University of Pittsburgh clinic and from cell lines from IDDM patients available from the Human Genetic Mutant Cell Repository, Camden, NJ) and non-diabetic controls (homozygous typing cells) using cloning methods. In one such method, which is a standard method, human genomic DNA was isolated from the patient samples using essentially the method of Maniatis et al., Molecular Cloning (1982), 280-281 or prepared from the buffy coat fraction, which is composed primarily of peripheral blood lymphocytes, as described by Saiki et al., Biotechnol. 3:1008-1012 (1985). This DNA was then cloned as full genomic libraries into bacteriophage vectors, as described in Maniatis, supra, pp. 269-294. Individual clones for the HLA-DR $\beta$  genes were selected by hybridization to radioactive cDNA probes (Maniatis, pp. 309-328) and characterized by restriction mapping. See U.S. Patent No. 4,582,788 issued April 15, 1986. Individual clones from IDDM patients were assigned to DR-typed haplotypes by comparing the done restriction map with the RFLP segregation pattern within the patient's family. Finally, small fragments of these clones representing the variable second exon were subcloned (Maniatis, pp. 390-402) into the Me3mp10 cloning vector, which is publicly available from Boehringer-Mannheim.

In an alternative procedure for cloning the genes, amplification of the relevant portion (the second exon) of the gene was carried out from a total of 1 microgram of each isolated human genomic DNA as described in Example I using primers GH46 and GH50, which have non-homologous sequences to act as linker/primers.

The reaction mixtures were subjected to 28 cycles of amplification and then the mixtures were stored at -20°C. Then the following cloning procedure was used for the amplified products.

The reaction mixture was sub-cloned into M13mp10 by first digesting in 50  $\mu$ l of buffer containing 50 mM NaCl, 10 mM Tris-HCl, pH 7.8, 10 mM MgCl<sub>2</sub>, 20 units PstI and 26 units HindIII at 37°C for 90 minutes. The reaction was stopped by freezing. The volume was adjusted to 110  $\mu$ l with a buffer containing Tris-HCl and EDTA and loaded onto a 1 ml BioGel P-4 spin dialysis column. One fraction was collected and ethanol precipitated.

The ethanol pellet was resuspended in 15  $\mu$ l water and adjusted to 20  $\mu$ l volume containing 50 mM Tris-HCl, pH 7.8, 10 mM MgCl<sub>2</sub>, 0.5 mM ATP, 10 mM dithiothreitol, 0.5  $\mu$ g M13mp10 vector digested with PstI and HindIII and 400 units ligase. This mixture was incubated for three hours at 16°C.

Ten microliters of ligation reaction mixture containing Molt 4 DNA was transformed into E. coli strain JM103 competent cells, which are publicly available from BRL in Bethesda, MD. The procedure followed for preparing the transformed strain is described in Messing, J. (1981) Third Cleveland Symposium on Macromolecules: Recombinant DNA, ed. A. Walton, Elsevier, Amsterdam, 143-153.

Eighteen of the alleles from these two cloning procedures were sequenced. In some of the sequences determined four areas of specific DNA and protein sequence were found to occur in various combinations and to be associated with IDDM. The DNA sequences seen in the genomes of IDDM patients produced an alteration in one to three amino acid residues of the DR $\beta$  protein. These four variable regions of the DR $\beta$  second exon are found in sequences obtained from many diabetic sources and are identified above. These regions can be used for synthesizing primers and probes used for detecting such sequences. These IDDM related sequences are identifiable as LR-S, -FL-, V-S, and I-DE, with LR-S and -FL- being found at amino acid residue positions 10 to 13 and 36 to 39 in the DR beta second exon.

### V. Primers and Amplification

Primers GH46 and GH50 described in Example CI may be employed to amplify DNA samples to be tested for IDDM. The amplification procedure of Example I or IIIA may be employed, using also 10  $\mu$ l DMSO.

### VI. Expected Synthesis of Probes

Two of four labeled DNA probes, designated GH54 (V-S), 5' CCTGTCGCCGAGTCCTGG) and GH78 (I-DE, 5' GACATCCTGGAAGACGAGAGA) may be employed. These probes may be synthesized as described for the primers and labeled as described above.

### VII. Expected Dot Blot Hybridizations

Using the dot blot method generally described in Example I, under stringent conditions, the probes are expected



to have reasonable specificity for the portions of the allele being detected in genomic DNA samples.

#### D. Amplification and Detection of HLA-DP $\alpha$ and DP $\beta$ Sequences

The known DP $\beta$  sequences, showing the type of polymorphism already known, are depicted in Figure 6 of Trowsdale et al., *Immunological Reviews*, No. 85 (1985), p 5-43, at page 16. Further polymorphisms may be identified. Primers for the conserved segments and probes to the variable segments of these genes can be designed similarly to what is described above.

The nucleotide sequence of DP $\alpha$ 1 alleles obtained from cDNA clones showing the type of polymorphism already known are depicted in Figure 4 of Trowsdale et al., *supra*, at page 12.

The detection and amplification of such sequences may be clinically useful in bonemarrow transplantations and in tissue typing.

#### EXAMPLE IV

Frozen Molt 4 cells (a T cell line homozygous for normal  $\beta$ -globin from Human Genetic Mutant Cell Repository, Camden, NJ as GM2219C), SC1 cells (a EBV-transformed B cell line homozygous for the sickle cell allele from ATCC, Rockville, MD as CRL8756) and GM2064 cells (control described above having no  $\beta$ -globin or  $\delta$ -globin sequence) were thawed and resuspended in phosphate buffered saline, such that 10  $\mu$ l of cells containing cell numbers varying from 37 to 1200 for each type of cell line was obtained. Each cell line was mixed with 35  $\mu$ l water and then overlaid with mineral oil. The resulting suspension was heated at 95°C for 10 minutes. Then a total of 55  $\mu$ l of the reagents used to amplify the cell lines in Example I, including primers PC03 and PC04, was added.

The amplification procedure of Example I can then be used followed by the dot blot procedure. This direct use of the cells eliminates isolating the genomic DNA from the cell line or clinical sample.

#### EXAMPLE V

The procedure of Example I was used to amplify the genomic DNA from a known normal individual, a known sickle cell individual, and an individual with no  $\beta$ -globin gene sequences (GM2064), except that the amplification was automated as described in Example IIIA.

Three labeled DNA probes, designated RS31, RS32, and RS33, of the following sequences were prepared as follows:

5' -\*TCCTGAGGAGAAGTCTG-3' (RS31)  
 5' -\*CCTGAGGAGAAGTCT-3' (RS32)  
 5' -\*CTGAGGAGAAGTC-3' (RS33)

where \* indicates the label. These probes are 17, 15, and 13 bases long, respectively, and are complementary to the normal  $\beta$ -globin allele ( $\beta^A$ ). The probes were synthesized and labeled as described in Example I.

Probes RS32, RS33, and RS18 were tested for specificity for non-amplified cloned normal and sickle-cell globin sequences using the procedure described in Example I, except that the hybridization temperature was reduced from 55°C to below 32°C. At hybridization temperatures below 55°C the RS18 (19-mer) did not show specificity unless the salt concentration was reduced. The two shorter probes showed excellent specificity at the higher salt content.

The probes RS31, RS32, RS33, and RS18 were tested against the amplified genomic DNA. Because of the conditions of temperature and salt concentration, the 19-mer showed no specificity at a hybridization temperature of below 32°C. All of the shorter mers did show specificity. These results clearly demonstrate that the shorter probes can be selective; and that the conditions of selectivity were less extreme than those needed for the 19-mer.

The 17-mer globin probe RS31 worked optimally when hybridized below 32°C in 6 x SSPE (with 10 x Denhardt's and 0.1% SDS) and then washed in 0.1 x SSPE for 10 minutes at 42°C.

The 15-mer globin probe RS32 worked optimally when hybridized below 32°C in 6 x SSPE (with 10 x Denhardt's and 0.1% SDS) and then washed in 0.1 x SSPE for 10 minutes at 32°C.

The 13-mer globin probe RS33 worked optimally when hybridized below 32°C in 6 x SSPE (with 10 x Denhardt's and 0.1% SDS) and then washed in 0.1 x SSPE for 10 minutes at 25°C.

EXAMPLE VI

## I. Synthesis of the Primers

Two primers identified below were synthesized by the method described in Example I to amplify a portion of the second exon of the  $\beta$ -globin gene:

5'-ATTTTCCCACCCTTAGGCTG-3' (RS40)  
5'-GCTCACTCAGTGTGGCAAAG-3' (RS42)

This primer pair defines a 198 base pair amplification product that includes the sites of three relatively common  $\beta$ -thalassemia mutations-the codon 39 non sense mutation, the codon 41-42 frameshift deletion, and the codon 44 frameshift deletion.

Because the  $\beta$ -globin gene in the region of codons 39 to 42 is exactly homologous to delta-globin, the primers were designed to be specific for and only amplify  $\beta$ -globin. The RS40 primer spans the first intron-second exon junction where there are six base pair mismatches with  $\delta$ -globin. RS42 is positioned over codons 84 to 91 and also contains six mismatches with  $\delta$ -globin. These mismatches are sufficient to prevent hybridization of the primers to the  $\delta$ -globin gene. After 20 cycles of amplification, the overall efficiency of these primers was approximately 80% and corresponded to a 130,000-fold amplification. As expected, the amplification product contained no detectable  $\delta$ -globin DNA.

## II. Isolation of Human Genomic DNA From Clinical Samples

Five genomic clinical DNA samples of various  $\beta$ -thalassemia genotypes were obtained from Drs. Alan Scott and Haig Kazazian (Johns Hopkins University, Baltimore, MD). These samples were JH1 (normal/39 non), JH2 (39 non/39 non), JH3 (normal/41 deletion), JH4 (17 non/41 del), and JH5 (39 non/44 deletion).

## III. Amplification Reaction

One microgram portions of each DNA sample and of Molt 4 as control were diluted into a 100  $\mu$ l volume with 50 mM NaCl, 10 mM Tris-HCl (pH 7.6), 10 mM  $MgCl_2$ , 1  $\mu$ M primer PCO3, 1  $\mu$ M primer PCO4, 10% DMSO (v/v), 1.5 mM dATP, 1.5 mM dCTP, 1.5 mM dGTP, and 1.5 mM dTTP and subjected to 25 cycles of automated amplification as described in Example IIIA, adding 1 unit of Klenow fragment at each cycle.

## IV. Oligodeoxyribonucleotide Probes

Four DNA probes, designed XX1, XX2, XX3 and XX4 below, were provided by Drs. Scott and Kazazian at Johns Hopkins University:

5'-\*CCTTGGACCTAGAGGTTCT-3' (XX1)  
5'-\*CCTTGGACCCAGAGGTTCT-3' (XX2)  
5'-\*GGTTCTTTGAGTCCTTTGG-3' (XX3)  
5'-\*GGTT---GAGTCCTTTGGGGAT-3' (XX4)

where \* indicates the label later attached. The XX1 and XX2 pair was used to detect the codon 39 non-sense mutation, with XX1 complementary to the normal allele and XX2 complementary to the non sense mutant. The other pair of probes was designed to test for the 4142 frameshift deletion, with XX3 annealing to the normal allele and XX4 to the deletion mutant. Each of the probes was phosphorylated as described in Example I.

## V. Dot Blot Hybridizations

Four replicate dot blots were prepared, each spot containing one-eighteenth of the amplification product (56 ng of genomic DNA). Each filter was individually prehybridized in 8 ml 5 x SSPE, 5 x Denhardt's solution, 0.5% SDS for 15 minutes at 55°C. One-half pmole of each labeled probe (specific activities ranged from 1.8 to 0.7  $\mu$ Ci/pmole) was added and the hybridization was continued for an additional 60 minutes at the same temperature. The filters were washed twice at room temperature in 2 x sodium saline phosphate EDTA (SSPE), 0.1% SDS, for 5-10 minutes per wash,

followed by a high-stringency wash in 5 x SSPE, 0.1% SDS for 10 minutes at 60°C. Autoradiograms were developed after overnight and two-hour exposures with a single intensification screen.

#### VI. Autoradiogram Results

The results were consistent with the listed genotypes of each DNA sample. Each probe annealed only to those genomic sequences with which it was perfectly matched.

#### EXAMPLE VII

The method herein may also be applied for forensic uses, by amplifying a random polymorphic region, e.g., HLA or mitochondrial DNA, to detect, e.g., nucleic acids in any body samples such as, e.g., hair samples, semen and blood samples; and other samples containing DNA. The nucleic acid may be extracted from the sample by any means, and primers and probes are selected based on identifying characteristics or known characteristics of the nucleic acid being detected.

#### EXAMPLE VIII

##### I. Synthesis of the Primers

The primers PC03 and PC04 described in Example I were employed herein.

##### II. Isolation of Human Genomic DNA from Cell Line

High molecular weight genomic DNA was isolated from aT cell line, Molt 4, homozygous for normal  $\beta$ -globin available from the Human Genetic Mutant Cell Depository, Camden, NJ as GM2219C using essentially the method of Maniatis et al., Molecular Cloning (1982), 280-281.

##### III. Purification of a Polymerase From *Thermus aquaticus*

*Thermus aquaticus* strain YT1, available without restriction from the American Type Culture Collection, 12301 Paridawn Drive, Rockville, MD, as ATCC No. 25,104 was grown in flasks in the following medium:

Sodium Citrate	1 mM
Potassium Phosphate, pH 7.9	5 mM
Ammonium Chloride	10mM
Magnesium Sulfate	0.2 mM
Calcium Chloride	0.1 mM
Sodium Chloride	1 g/l
Yeast Extract	1 g/l
Tryptone	1 g/l
Glucose	2 g/l
Ferrous Sulfate	0.01 mM
(The pH was adjusted to 8.0 prior to autoclaving.)	

A 10-liter fermentor was inoculated from a seed flask cultured overnight in the above medium at 70°C. A total of 600 ml from the seed flask was used to inoculate 10 liters of the same medium. The pH was controlled at 8.0 with ammonium hydroxide with the dissolved oxygen at 40%, with the temperature at 70°C, and with the stirring rate at 400 rpm.

After growth of the cells, they were purified using the protocol (with slight modification) of Kaledin et al., supra, through the first five stages and using a different protocol for the sixth stage. All six steps were conducted at 4°C. The rate of fractionation on columns was 0.5 column volumes/hour and the volumes of gradients during elution were 10 column volumes.

Briefly, the above culture of the *T. aquaticus* cells was harvested by centrifugation after nine hours of cultivation, in late log phase, at a cell density of 1.4 g dry weight/l. Twenty grams of cells was resuspended in 80 ml of a buffer consisting of 50 mM Tris-HCl pH 7.5, 0.1 mM EDTA. Cells were lysed and the lysate was centrifuged for two hours at

35,000 rpm in a Beckman TI 45 rotor at 4°C. The supernatant was collected (fraction A) and the protein fraction precipitating between 45 and 75% saturation of ammonium sulfate was collected, dissolved in a buffer consisting of 0.2 M potassium phosphate buffer, pH 6.5, 10 mM 2-mercaptoethanol, and 5% glycerine, and finally dialyzed against the same buffer to yield fraction B.

Fraction B was applied to a 2.2 x 30-cm column of DEAE-cellulose, equilibrated with the above described buffer. The column was then washed with the same buffer and the fractions containing protein (determined by absorbance at 280 nm) were collected. The combined protein fraction was dialyzed against a second buffer, containing 0.01 M potassium phosphate buffer, pH 7.5, 10 mM 2-mercaptoethanol, and 5% glycerine, to yield fraction C.

Fraction C was applied to a 2.6 x 21-cm column of hydroxyapatite, equilibrated with a second buffer. The column was then washed and the enzyme was eluted with a linear gradient of 0.01-0.5 M potassium phosphate buffer, pH 7.5, containing 10 mM 2-mercaptoethanol and 5% glycerine. Fractions containing DNA polymerase activity (90-180 mM potassium phosphate) were combined, concentrated four-fold using an Amicon stirred cell and YM10 membrane, and dialyzed against the second buffer to yield fraction D.

Fraction D was applied to a 1.6 x 28-cm column of DEAE-cellulose, equilibrated with the second buffer. The column was washed and the polymerase was eluted with a linear gradient of 0.01-0.5 M potassium phosphate in the second buffer. The fractions were assayed for contaminating endonuclease(s) and exonuclease(s) by electrophoretically detecting the change in molecular weight of phage  $\lambda$  DNA or supercoiled plasma DNA after incubation with an excess of DNA polymerase (for endonuclease) and after treatment with a restriction enzyme that cleaves the DNA into several fragments (for exonuclease). Only those DNA polymerase fractions (65-95 mM potassium phosphate) having minimal nuclease contamination were pooled. To the pool was added autoclaved gelatin in an amount of 250  $\mu$ g/ml, and dialysis was conducted against the second buffer to yield Fraction E.

Fraction E was applied to a 9 ml phosphocellulose column and eluted with a 100 ml gradient (0.01-0.4 M KCl gradient in 20 mM potassium phosphate buffer pH 7.5). The fractions were assayed for contaminating endo/exonuclease(s) as described above as well as for polymerase activity (by the method of Kaledin et al.) and then pooled. The pooled fractions were dialyzed against the second buffer, then concentrated by dialysis against 50% glycerine and the second buffer.

The molecular weight of the polymerase was determined by SDS PAGE. Marker proteins (Bio-Rad low molecular weight standards) were phosphorylase B (92,500), bovine serum albumin (66,200), ovalbumin (45,000), carbonic anhydrase (31,000), soybean trypsin inhibitor (21,500), and lysozyme (14,400).

Preliminary data suggest that the polymerase has a molecular weight of about 86,000-90,000 daltons, not 62,000-63,000 daltons reported in the literature (e.g., by Kaledin et al.).

#### IV. Amplification Reaction

One microgram of the genomic DNA described above was diluted in an initial 100  $\mu$ l aqueous reaction volume containing 25 mM Tris-HCl buffer (pH 8.0), 50 mM KCl, 10 mM  $MgO_2$ , 5 mM dithiothreitol, 200  $\mu$ g/ml gelatin, 1  $\mu$ M of primer PC03, 1  $\mu$ M of primer PC04, 1.5 mM dATP, 1.5 mM dCTP, 1.5 mM dGTP and 1.5 mM TTP. The sample was heated for 10 minutes at 98°C to denature the genomic DNA, then cooled to room temperature. Four microliters of the polymerase from *Thermus aquaticus* was added to the reaction mixture and overlaid with a 100  $\mu$ l mineral oil cap. The sample was then placed in the aluminum heating block of the liquid handling and heating instrument described in copending U.S. Application Serial No. 833,368 filed February 25, 1986, the disclosure of which is incorporated herein by reference.

The DNA sample underwent 20 cycles of amplification in the machine, repeating the following program cycle:


- 1) heating from 37°C to 98°C in heating block over a period of 2.5 minutes; and
- 2) cooling from 98°C to 37°C over a period of three minutes to allow the primers and DNA to anneal. After the last cycle, the sample was incubated for an additional 10 minutes at 55°C to complete the final extension reaction.

#### V. Synthesis and Phosphorylation of Ologodeoxynbonucleotide Probes

A labeled DNA probe, designated RS24, of the following sequence was prepared:

**5' -\*CCCACAGGGCAGTAACGGCAGACTTCTCCTCAGGAGTCAG-3' (RS24)**

where \* indicates the label. This probe is 40 bases long, spans the fourth through seventeenth codons of the gene, and is complementary to the normal  $\beta$ -globin allele ( $\beta^A$ ). The schematic diagram of primers and probes is given below:



This probe was synthesized according to the procedures described in Section I of Example I. The probe was labeled by contacting 20 pmole thereof with 4 units of T4 polynucleotide kinase and about 40 pmole  $\gamma$ - $^{32}\text{P}$ -ATP (about 7000 Ci/mmole) in a 40  $\mu\text{l}$  reaction volume containing 70 mM Tris buffer (pH 7.6), 10 mM  $\text{MgCl}_2$ , 1.5 mM spermine and 10 mM dithiothreitol for 60 minutes at 37°C. The total volume was then adjusted to 100  $\mu\text{l}$  with 25 mM EDTA and purified according to the procedure of Maniatis et al., *Molecular Cloning* (1982), 466-467 over a 1 ml spin dialysis column equilibrated with Tris-EDTA (TE) buffer (10 mM Tris buffer, 0.1 mM EDTA, pH 8.0). TCA precipitation of the reaction product indicated that for RS24 the specific activity was 4.3  $\mu\text{Ci}/\text{pmole}$  and the final concentration was 0.118 pmole/ $\mu\text{l}$ .

#### VI. Dot Blot Hybridizations

Four microliters of the amplified sample from Section 1 and 5.6  $\mu\text{l}$  of appropriate dilutions of  $\beta$ -globin plasmid DNA calculated to represent amplification efficiencies of 70, 75, 80, 85, 90, 95 and 100% were diluted with 200  $\mu\text{l}$  0.4 N NaOH, 25 mM EDTA and spotted onto a nylon filter by first wetting the filter with water, placing it in an apparatus for preparing dot blots which holds the filters in place, applying the samples, and rinsing each well with 0.1 ml of 20 x SSPE (3.6 M NaCl, 200 mM  $\text{NaH}_2\text{PO}_4$ , 20 mM EDTA), as disclosed by Reed and Mann, *Nucleic Acids Research*, 13, 7202-7221 (1985). The filters were then removed, rinsed in 20 x SSPE, and baked for 30 minutes at 80°C in a vacuum oven.

After baking, each filter was then contacted with 16 ml of a hybridization solution consisting of 3 x SSPE, 5 x Denhardt's solution (1 x = 0.02% polyvinylpyrrolidone, 0.02% Ficoll, 0.02% bovine serum albumin, 0.2 mM Tris, 0.2 mM EDTA, pH 8.0), 0.5% SDS, and 30% formamide, and incubated for two hours at 42°C. Then 2 pmole of probe RS24 was added to the hybridization solution and the filter was incubated for two hours at 42°C.

Finally, each hybridized filter was washed twice with 100 ml of 2 x SSPE and 0.1% SDS for 10 minutes at room temperature. Then the filters were treated once with 100 ml of 2 x SSPE, 0.1% SDS at 60°C for 10 minutes.

Each filter was then autoradiographed, with the signal readily apparent after two hours.

#### VII. Discussion of Autoradiogram

The autoradiogram of the dot blots was analyzed after two hours and compared in intensity to standard serial dilution  $\beta$ -globin reconstructions prepared with *Hae*III/*Mae*I-digested pBR:  $\beta^A$ , where  $\beta^A$  is the wild-type allele, as described in Saiki et al., *Science*, *supra*. Analysis of the reaction product indicated that the overall amplification efficiency was about 95%, corresponding to a 630,000-fold increase in the  $\beta$ -globin target sequence.

### EXAMPLE IX

#### I. Amplification Reaction

Two 1  $\mu\text{g}$  samples of genomic DNA extracted from the Molt 4 cell line as described in Example VIII were each diluted in a 100  $\mu\text{l}$  reaction volume containing 50 mM KCl, 25 mM Tris-HCl buffer pH 8.0, mM  $\text{MgCl}_2$ , 1  $\mu\text{M}$  of primer PC03, 1  $\mu\text{M}$  of primer PC04, 200  $\mu\text{g}/\text{ml}$  gelatin, 10% dimethylsulfoxide (by volume), 1.5 mM dATP, 1.5 mM dCTP, 1.5 mM dGTP, and 1.5 mM TTP. After this mixture was heated for 10 minutes at 98°C to denature the genomic DNA, the samples were cooled to room temperature and 4  $\mu\text{l}$  of the polymerase from *Thermus aquaticus* described in Example VIII was added to each sample. The samples were overlaid with mineral oil to prevent condensation and evaporative loss.

One of the samples was placed in the heating block of the machine described in Example VIII and subjected to 25 cycles of amplification, repeating the following program cycle:

- (1) heating from 37 to 93°C over a period of 2.5 minutes;
- (2) cooling from 93°C to 37°C over a period of three minutes to allow the primers and DNA to anneal; and
- (3) maintaining at 37°C for two minutes.

After the last cycle the sample was incubated for an additional 10 minutes at 60°C to complete the final extension.

reaction.

The second sample was placed in the heat-conducting container of a machine, which container is attached to Peltier heat pumps which adjust the temperature upwards or downwards and a microprocessor controller to control automatically the amplification sequence, the temperature levels, the temperature ramping and the timing of the temperature.

The second sample was subjected to 25 cycles of amplification, repeating the following program cycle:

- (1) heating from 37 to 95°C over a period of three minutes;
- (2) maintaining at 95°C for 0.5 minutes to allow denaturation to occur;
- (3) cooling from 95 to 37°C over a period of one minute; and
- (4) maintaining at 37°C for one minute.


## II. Analysis

Two tests were done for analysis, a dot blot and an agarose gel analysis.

For the dot blot analysis, a labeled DNA probe, designated RS18, of the following sequence was prepared.

5'--\*CTCCTGAGGAGAAGTCTGC-3' (RS18)

where \* indicates the label. This probe is 19 bases long, spans the fourth through seventeenth codons of the gene, and is complementary to the normal  $\beta$ -globin allele ( $\beta^A$ ). The schematic diagram of primers and probes is given below:



This probe was synthesized according to the procedures described in Section I of Example I. The probe was labeled by contacting 10 pmole thereof with 4 units of T4 polynucleotide kinase and about 40 pmole  $\gamma$ - $^{32}\text{P}$ -ATP (about 7000 Ci/mmole) in a 40  $\mu\text{l}$  reaction volume containing 70 mM Tris-HCl buffer (pH 7.6), 10 mM  $\text{MgCl}_2$ , 1.5 mM spermine and 10 mM dithiothreitol for 60 minutes at 37°C. The total volume was then adjusted to 100  $\mu\text{l}$  with 25 mM EDTA and purified according to the procedure of Maniatis et al., *Molecular cloning* (1982), 466-467 over a 1 ml spin dialysis column equilibrated with Tris-EDTA (TE) buffer (10 mM Tris-HCl buffer, 0.1 mM EDTA, pH 8.0). TCA precipitation of the reaction product indicated that for RS18 the specific activity was 4.6  $\mu\text{Ci}/\text{pmole}$  and the final concentration was 0.114  $\text{pmole}/\mu\text{l}$ .

Five microliters of the amplified sample from Section I and of a sample amplified as described above except using the Klenow fragment of *E. coli* DNA Polymerase I instead of the thermostable enzyme were diluted with 195  $\mu\text{l}$  0.4 N NaOH, 25 mM EDTA and spotted onto two replicate cationic nylon filters by first wetting the filters with water, placing them in an apparatus for preparing dot blots which holds the filters in place, applying the samples, and rinsing each well with 0.4 ml of 20 x SSPE (3.6 M NaCl, 200 mM  $\text{NaH}_2\text{PO}_4$ , 20 mM EDTA), as disclosed by Reed and Mann, *Nucleic Acids Research*, 13, 7202-7221 (1985). The filters were then removed, rinsed in 20 x SSPE, and baked for 30 minutes at 80°C in a vacuum oven.

After baking, each filter was then contacted with 6 ml of a hybridization solution consisting of 5 x SSPE, 5 x Denhardt's solution (1 x = 0.02% polyvinylpyrrolidone, 0.02% Ficoll, 0.02% bovine serum albumin, 0.2 mM Tris, 0.2 mM EDTA, pH 8.0) and 0.5% SDS, and incubated for 60 minutes at 55°C. Then 5  $\mu\text{l}$  of probe RS18 was added to the hybridization solution and the filter was incubated for 60 minutes at 55°C.

Finally, each hybridized filter was washed twice with 100 ml of 2 x SSPE and 0.1% SDS for 10 minutes at room temperature. Then the filters were treated twice more with 100 ml of 5 x SSPE, 0.1% SDS at 60°C for 1) one minute and 2) three minutes, respectively.

Each filter was then autoradiographed, with the signal readily apparent after 90 minutes.

In the agarose gel analysis, 5  $\mu\text{l}$  each amplification reaction was loaded onto 4% NuSieve 0.5% agarose gel in 1 x TBE buffer (0.089 M Tris borate, 0.089 M boric acid, and 2 mM EDTA) and electrophoresed for 60 minutes at 100V. After staining with ethidium bromide, DNA was visualized by UV fluorescence.

The results show that the machines used in Examples VIII and IX herein were equally effective in amplifying the DNA, showing discrete high-intensity 110-base pair bands of similar intensity, corresponding to the desired sequence,

as well as a few other discrete bands of much lower intensity. In contrast, the amplification method, which involves reagent transfer after each cycle using the Klenow fragment of *E. coli* Polymerase I, gave a DNA smear resulting from the non-specific amplification of many unrelated DNA sequences.

It is expected that similar improvements in amplification and detection would be achieved in evaluating HLA-DQ, DR and DP regions.

It is also expected that the procedure of Example I may be repeated using a biotinylated probe prepared as described in U.S. Patent Nos. 4,582,789 and 4,617,261.

#### EXAMPLE X

cDNA was made from 1 µg of rabbit reticulocyte mRNA in a 100 µl reaction volume containing 150 mM KCl, 50 mM Tris-HCl (pH 8.3), 10 mM MgCl<sub>2</sub>, 5 mM DTT, 0.5 mM dATP, 0.5 mM dCTP, 0.5 mM dTTP, 0.5 mM dGTP, 0.2 µg oligo(dT)<sub>12-18</sub>, 40 units RNasin, and 5 units AMV reverse transcriptase, and incubated for 30 minutes at 42°C. The reaction was stopped by heating for 10 minutes at 95°C. Two µg RNase A was added to the sample (2 µl of a 2 mg/ml solution in water) and incubated for 10 minutes at 37°C.

Three amplification reactions were done with the Klenow fragment using different pairs of primers. The primer pair PC03/PC04 define a 110-bp product the primer pair RS45/oligo(dT)<sub>25-30</sub> define an about 370-bp product, and the primer pair PC03/oligo(dT)<sub>25-30</sub> an about 600-bp product. PC03, PC04, and RS45 are complementary to the human β-globin gene and each has two mismatches with the rabbit gene. PC03 and PC04 are described in Example I. RS45 has the sequence: 5'-CAAGAAGGTGCTAGGTGCC-3'.

The amplification reactions were performed with 1/20th (5 µl) of the cDNA described above in a 100 µl reaction volume containing 50 mM NaCl, 10 mM Tris-HCl (pH 7.6), 10 mM MgCl<sub>2</sub>, 200 µg/ml gelatin, 10% DMSO, 1 µM PC03 or RS45, 1 µM PC04 or oligo(dT)<sub>25-30</sub>, 1.5 mM dATP, 1.5 mM dCTP, 1.5 mM dTTP and 1.5 mM dGTP. The samples were heated for five minutes at 98°C, then cooled to room temperature and overlaid with about 100 µl mineral oil.

The samples were subjected to 10 cycles of automated amplification using the machine described in Example VIII and using the following program:

- 1) heating from 37°C to 98°C in heating block over 2.5 minutes (denature);
- 2) cooling from 98°C to 37°C over 3.0 minutes (anneal);
- 3) adding 1 unit Klenow fragment; and
- 4) maintaining at 37°C for 20 minutes (extend).

The final volume of each sample was about 140 µl.

One-twentieth (7 µl) of each sample was analyzed by electrophoresis on a 2% agarose gel. After staining with ethidium bromide, discrete bands were seen in the PC03/PC04 and RS45/oligo(dT) samples. The sizes of the bands were consistent with the expected lengths: 110-bp for the former, about 370-bp for the latter. No evidence of amplification of an about 600-bp fragment with the PC03/oligo(dT) primer pair was observed.

The contents of the gel were Southern blotted onto a cationic nylon membrane and hybridized with a nick-translated human β-globin probe, pBR328:β<sub>g</sub>αA, described in Saiki et al., *Science*, *supra*, using standard techniques. The resulting autoradiogram extended the conclusions reached previously - the 110 and about 370-bp fragments were β-globin specific amplification products and no significant amplification of the about 600-bp band was detected.

Three additional samples were amplified with the *Thermus aquaticus* (Taq) polymerase obtained as described above using the same primer pairs described previously. Five microliter portions of cDNA were amplified in 100 µl reaction volumes containing 50 mM KCl, 25 mM Tris-HCl (pH 8.0), 10 mM MgCl<sub>2</sub>, 200 µg/ml gelatin, 10% DMSO, 1 µM PC03 or RS45, 1 µM PC04 or oligo(dT)<sub>25-30</sub>, 1.5 mM dATP, 1.5 mM dCTP, 1.5 mM dTTP and 1.5 mM dGTP. The samples were heated for five minutes at 98°C, then cooled to room temperature. One microliter of Taq polymerase (1/8 dilution of lot 2) was added to each and overlaid with about 100 µl mineral oil.

The samples were subjected to 9 cycles of amplification in the Peltier device described in the previous example using the following program:

- 1) 1 min, 35 to 60°C ramp;
- 2) 12 min, 60 to 70°C ramp (extend);
- 3) 1 min, 70 to 95°C ramp (denature);
- 4) 30 sec, 95°C soak;
- 5) 1 min, 95 to 35°C ramp (anneal); and
- 6) 30 sec, 35°C soak.

After the last cycle, the samples were incubated an additional 10 minutes at 60°C to complete the final (10th cycle)

extension. The final volume of each was about 100  $\mu$ l.

As before, 1/20th (10  $\mu$ l) of each sample was analyzed on a 2% agarose gel. In this gel, amplification products were present in all three samples: 110-bp for PC03/PC04, about 370-bp for RS45/oligo(dT), and about 600-bp for PC03/oligo(dT). These results were confirmed by Southern transfer and hybridization with the pBR-328:betaA probe.

The production of the 600-bp product with Taq polymerase but not with the Klenow fragment is significant, and suggests that Taq polymerase is capable of producing longer DNA than the Klenow fragment. In summary, the technique herein wherein nucleic acids are amplified in a chain reaction in which primer extension products are produced which can subsequently act as templates, and the amplified samples are analyzed using sequence-specific probes provides several important advantages. It is a simplified procedure because the amplified samples can be spotted on a filter membrane as a dot blot, thereby avoiding the restriction digestion, electrophoresis and gel manipulations otherwise required. It is a more specific procedure because the amplification greatly increases the ratio of specific target sequence to cross-hybridizing sequences.

In addition, the process herein improves sensitivity by  $10^3$ - $10^4$ . An interpretable signal can be obtained with a 1 ng sample after an overnight exposure. Finally, by increasing the amount of sample applied to the filter to 0.1 to 0.5  $\mu$ g, it is possible that biotinylated oligonucleotide probes may be utilized.

#### Claims

1. A process for detecting the presence of a specific nucleotide sequence in nucleic acid in a sample, which process comprises:

(a) treating the sample, together or sequentially, with four different nucleoside triphosphates, an agent for polymerization of the nucleoside triphosphates, and two oligonucleotide primers for said nucleic acid under hybridizing conditions such that a primer will hybridize to said nucleic acid and an extension product of the primer be synthesized which is complementary to said nucleic acid, wherein said primers are selected such that the extension product synthesized from one primer, when separated from its complement, can serve as a template for synthesis of the extension product of the other primer;

(b) treating the sample under denaturing conditions to separate the primer extension products from their templates;

(c) treating the sample, together or sequentially, with said four nucleoside triphosphates, an agent for polymerization of the nucleoside triphosphates, and oligonucleotide primers such that a primer extension product is synthesized using each of the single strands produced in step (b) as a template, wherein steps (b) and (c) are repeated a sufficient number of times exponentially to increase the amount of said nucleic acid and to result in detectable amplification thereof;

(d) directly transferring, without gel fractionation, product derived from step (c) to a membrane;

(e) treating the membrane from (d) under hybridization conditions with a labeled sequence-specific oligonucleotide probe capable of hybridizing with the amplified nucleic acid only if the sequence of the probe is complementary to a region of the amplified nucleic acid; and

(f) detecting whether the probe has hybridized to an amplified nucleic acid in the sample.

2. A process of Claim 1, wherein the nucleic acid in the sample is extracted from the sample prior to step (a), and steps (b) and (c) are repeated at least five times, optionally wherein the sample contains human genomic DNA and steps (b) and (c) are repeated 15-30 times.

3. A process of Claim 1 or Claim 2, wherein said nucleic acid is double-stranded and its strands are separated by denaturing before or during step (a).

4. A process of Claim 1 or Claim 2, wherein the said nucleic acid is single-stranded.

5. A process of Claim 3, wherein said nucleic acid is DNA and said primers are oligodeoxyribonucleotides.

6. A process of Claim 4, wherein said nucleic acid is DNA or messenger RNA and said primers are oligodeoxyribonucleotides.

7. A process of any one of Claims 1 to 6, wherein said agent for polymerization is selected from *E. coli* DNA Polymerase I, Klenow fragment of *E. coli* DNA Polymerase I, T4 DNA polymerase, thermostable polymerase enzyme and reverse transcriptase.



8. A process of any one of Claims 1 to 7, wherein the specific nucleotide sequence is contained within a human nucleic acid selected from a beta-globin gene, an oncogene, an HLA-DQ alpha, DO beta, DR beta, DP alpha and DP beta gene, and mitochondrial DNA, optionally wherein the nucleotide sequence is associated with insulin-dependent diabetes mellitus or wherein the nucleotide sequence is contained within nucleic acid of an organism that can cause an infectious disease.

9. A process of any one of Claims 1 to 8, wherein either the sample comprises cells, optionally wherein the sample is peripheral blood lymphocytes or amniotic fluid, and the process further comprises, before step (a), the step of heating the sample sufficiently to expose the nucleic acid(s) therein; or wherein the sample is a semen, hair or blood sample, or contains mitochondrial DNA, and the process is used in forensic analysis.

10. A process for detecting the presence of a specific nucleotide sequence in a nucleic acid in a sample, which process comprises:

(a) treating the sample, together or sequentially, with four different nucleoside triphosphates, an agent for polymerization of the nucleoside triphosphates, and two oligonucleotide primers for said nucleic acid under hybridizing conditions such that a primer hybridizes to said nucleic acid and an extension product of the primer is synthesized which is complementary to said nucleic acid, wherein said primers are selected such that the extension product synthesized from one primer, when separated from its complement, can serve as a template for synthesis of the extension product of the other primer;

(b) treating the sample under denaturing conditions to separate the primer extension products from their templates;

(c) treating the sample, together or sequentially, with said four nucleoside triphosphates, an agent for polymerization of the nucleoside triphosphates, and oligonucleotide primers such that a primer extension product is synthesized using each of the single strands produced in step (b) as a template, wherein steps (b) and (c) are repeated a sufficient number of times exponentially to increase the amount of said nucleic acid and to result in detectable amplification thereof; and wherein at least one primer and/or at least one of the four nucleoside triphosphates is labeled with a detectable moiety;

(d) treating a membrane to which is attached a sequence-specific oligonucleotide capable of hybridizing with the amplified nucleic acid only if the sequence of the oligonucleotide is complementary to a region of the amplified nucleic acid under hybridization conditions with product derived from step (c); and

(e) detecting whether an amplified nucleic acid in the sample has hybridized to the oligonucleotide attached to the membrane.

11. A process for detecting the presence of a specific nucleotide sequence in nucleic acid in a sample, which process comprises:

(a) treating the sample, together or sequentially, with four different nucleoside triphosphates, an agent for polymerization of the nucleoside triphosphates, and two oligonucleotide primers for said nucleic acid under hybridizing conditions such that a primer hybridizes to said nucleic acid and an extension product of the primer is synthesized which is complementary to said nucleic acid, wherein said primers are selected such that the extension product synthesized from one primer, when separated from its complement, can serve as a template for synthesis of the extension product of the other primer;

(b) treating the sample under denaturing conditions to separate the primer extension products from their templates;

(c) treating the sample, together or sequentially, with said four nucleoside triphosphates, an agent for polymerization of the nucleoside triphosphates, and oligonucleotide primers such that a primer extension product is synthesized using each of the single strands produced in step (b) as a template, wherein steps (b) and (c) are repeated a sufficient number of times exponentially to increase the amount of said nucleic acid and to result in detectable amplification thereof;

(d) attaching to a membrane a labeled sequence-specific oligonucleotide probe capable of hybridizing with the amplified nucleic acid only if the sequence of the probe is complementary to a region of the amplified nucleic acid;

(e) treating the membrane from (d) under hybridization conditions with product derived from step (c);

(f) treating the product of step (e) with a restriction enzyme which will cleave any hybrids formed if the probe and any sequence to be detected both contain a restriction site recognized by the enzyme; and

(g) detecting whether a labeled restriction fragment of the required length is present in the restriction digest thereby indicating that hybridization occurred.

12. A process of Claim 10, wherein said primers are oligodeoxyribonucleotides and are optionally labeled and said agent for polymerization is a DNA polymerase.

13. A process of Claim 10 or Claim 12, wherein said nucleic acid is DNA, optionally human genomic DNA.

14. A process of Claim 10 or Claim 12, wherein said nucleic acid is RNA.

15. A process of Claim 13, wherein said primers amplify a segment of a Class II HLA gene, optionally a DQ alpha allele, a DQ beta allele, a DP alpha allele, a DP beta allele, a DR alpha allele, or a DR beta allele.

16. A kit for detecting the presence of a specific nucleotide sequence in a nucleic acid in a sample, which kit comprises, in packaged form, a multicontainer unit having:

(a) two oligonucleotide primers for said nucleic acid, which primers are selected such that an extension product synthesized from one primer, when separated from its complement, can serve as a template for the synthesis of the extension product of the other primer so as exponentially to amplify said nucleic acid;

(b) an agent for polymerization;

(c) four different nucleoside triphosphates; and either:

(d) a sequence specific oligonucleotide probe capable of hybridizing with the amplified nucleic acid spotted directly on a membrane only if the sequence of the probe is complementary to a region of the amplified nucleic acid; and

(e) a membrane to which amplified nucleic acid can directly bind for subsequent detection by probe hybridization; or

(d) a sequence specific oligonucleotide probe attached to a membrane, wherein said probe is capable of hybridizing with the amplified nucleic acid only if the sequence of the probe is complementary to a region of the amplified nucleic acid.

17. A kit of Claim 16, wherein either said primers are oligodeoxyribonucleotides, said nucleoside triphosphates are dATP, dCTP, dGTP and TTP, and the probe is labeled or wherein at least one of the primers and/or at least one of the nucleoside triphosphates is labeled.

18. A kit of Claim 16 or Claim 17, further comprising means of separating the strands of any double-stranded nucleic acids contained in the sample, and/or a positive control for the probe which contains one or more nucleic acids with the specific nucleotide sequence to be detected and a negative control for the probe which does not contain nucleic acid with the nucleotide sequence to be detected.

19. A kit of any one of Claims 16 to 18, wherein the agent for polymerization is an enzyme selected from E coli DNA Polymerase I, Klenow fragment of E coli DNA Polymerase I, T4 DNA polymerase, a heat-stable DNA polymerase enzyme, and reverse transcriptase.

20. A kit of any one of Claims 16 to 19 wherein the specific nucleotide sequence is contained within a human nucleic acid selected from a beta-globin gene, an HLA-DQ alpha, DQ beta, DR beta, DP alpha and DP beta gene, and mitochondrial DNA, optionally the nucleotide variation in sequence being associated with insulin-dependent diabetes mellitus.

## Patentansprüche

1. Verfahren zum Nachweis des Vorhandenseins einer spezifischen Nukleotidsequenz in Nukleinsäure in einer Probe, bei welchem:

(a) die Probe gleichzeitig oder nacheinander mit vier verschiedenen Nukleosidtriphosphaten, einem Mittel zur Polymerisation der Nukleosidtriphosphate und zwei Oligonukleotidprimern für die Nukleinsäure unter hybridisierenden Bedingungen behandelt wird, so daß ein Primer an die Nukleinsäure hybridisieren wird und ein Extensionsprodukt des Primers synthetisiert wird, das komplementär zu der Nukleinsäure ist, wobei die Primer so ausgewählt sind, daß das mit einem Primer synthetisierte Extensionsprodukt als Matrize für die Synthese des Extensionsprodukts des anderen Primers dienen kann, wenn es von seinem Komplement getrennt wird;

(b) die Probe unter denaturierenden Bedingungen behandelt wird, um die Primerextensionsprodukte von ihren Matrizen abzutrennen;

(c) die Probe gleichzeitig oder nacheinander mit den vier Nukleosidtriphosphaten, einem Mittel zur Polymerisation der Nukleosidtriphosphate und Oligonukleotidprimern behandelt wird, so daß ein Primerextensionsprodukt unter Verwendung jeder der in Schritt (b) hergestellten Einzelstränge als Matrize synthetisiert wird, wobei die Schritte (b) und (c) ausreichend oft wiederholt werden, um exponentiell die Menge der Nukleinsäure zu erhöhen und zu einer nachweisbaren Amplifikation davon zu führen:

(d) das aus Schritt (c) erhaltene Produkt direkt, ohne Gelfraktionierung, auf eine Membran übertragen wird;

(e) die Membran aus (d) unter hybridisierenden Bedingungen mit einer markierten sequenzspezifischen Oligonukleotidsonde behandelt wird, die mit der amplifizierten Nukleinsäure nur dann hybridisiert, wenn die Sequenz der Sonde komplementär zu einer Region der amplifizierten Nukleinsäure ist; und

(f) nachgewiesen wird, ob die Sonde an eine amplifizierte Nukleinsäure in der Probe hybridisiert hat.

2. Verfahren nach Anspruch 1, wobei die Nukleinsäure in der Probe vor Schritt (a) aus der Probe extrahiert wird, und die Schritte (b) und (c) wenigstens fünfmal wiederholt werden, wobei gegebenenfalls die Probe menschliche genomische DNA enthält und die Schritte (b) und (c) 15 bis 30mal wiederholt werden.

3. Verfahren nach Anspruch 1 oder Anspruch 2, wobei die Nukleinsäure doppelsträngig ist und ihre Stränge vor oder während Schritt (a) durch Denaturieren getrennt werden.

4. Verfahren nach Anspruch 1 oder Anspruch 2, wobei die Nukleinsäure einzelsträngig ist.

5. Verfahren nach Anspruch 3, wobei die Nukleinsäure DNA ist und die Primer Oligodeoxyribonukleotide sind.

6. Verfahren nach Anspruch 4, wobei die Nukleinsäure DNA oder Messenger-RNA ist und die Primer Oligodeoxyribonukleotide sind.

7. Verfahren nach einem der Ansprüche 1 bis 6, wobei das Mittel zur Polymerisation ausgewählt wird aus E. coli-DNA-Polymerase I, Klenow-Fragment von E coli-DNA-Polymerase I, T4-DNA-Polymerase, thermostabilem Polymeraseenzym und reverser Transkriptase.

8. Verfahren nach einem der Ansprüche 1 bis 7, wobei die spezifische Nukleotidsequenz innerhalb einer menschlichen Nukleinsäure enthalten ist, ausgewählt aus einem beta-Globin, einem Oncogen, einem HLA-DQ-alpha-, -DQ-beta-, -DR-beta-, -DP-alpha- und -DP-beta-Gen, und mitochondrialer DNA, wobei gegebenenfalls die Nukleotidsequenz mit insulinabhängigem Diabetes mellitus assoziiert ist, oder wobei die Nukleotidsequenz innerhalb einer Nukleinsäure eines Organismus enthalten ist, der eine infektiöse Erkrankung verursachen kann.

9. Verfahren nach einem der Ansprüche 1 bis 8, wobei die Probe entweder Zellen enthält, wobei es sich gegebenenfalls bei der Probe um periphere Blutlymphozyten oder amniotische Flüssigkeit handelt, und das Verfahren vor Schritt (a) ferner aufweist, daß die Probe ausreichend erhitzt wird, um die Nukleinsäure(n) darin zu exponieren; oder wobei es sich bei der Probe um eine Sperma-, Haar- oder Blutprobe handelt, oder mitochondriale DNA enthält, und das Verfahren in der forensischen Analyse verwendet wird.

10. Verfahren zum Nachweis des Vorhandenseins einer spezifischen Nukleotidsequenz in einer Nukleinsäure in einer Probe, bei welchem:

(a) die Probe gleichzeitig oder nacheinander mit vier verschiedenen Nukleosidtriphosphaten, einem Mittel zur Polymerisation der Nukleosidtriphosphate und zwei Oligonukleotidprimern für die Nukleinsäure unter hybridisierenden Bedingungen behandelt wird, so daß ein Primer an die Nukleinsäure hybridisiert und ein Extensionsprodukt des Primers synthetisiert wird, das komplementär zu der Nukleinsäure ist, wobei die Primer so ausgewählt sind, daß das mit einem Primer synthetisierte Extensionsprodukt als Matrize für die Synthese des Extensionsprodukts des anderen Primers dienen kann, wenn es von seinem Komplement getrennt wird;

(b) die Probe unter denaturierenden Bedingungen behandelt wird, um die Primerextensionsprodukte von ihren

Matrizen abzutrennen;

(c) die Probe gleichzeitig oder nacheinander mit den vier Nukleosidtriphosphaten, einem Mittel zur Polymerisation der Nukleosidtriphosphate und Oligonukleotidprimern behandelt wird, so daß ein Primerextensionsprodukt unter Verwendung jeder der in Schritt (b) hergestellten Einzelstränge als Matrize synthetisiert wird, wobei die Schritte (b) und (c) ausreichend oft wiederholt werden, um exponentiell die Menge der Nukleinsäure zu erhöhen und zu einer nachweisbaren Amplifikation davon zu führen; und wobei mindestens ein Primer und/oder mindestens eines der vier Nukleosidtriphosphate mit einem nachweisbaren Rest markiert ist;

(d) eine Membran, an der ein sequenzspezifisches Oligonukleotid befestigt ist, das mit der amplifizierten Nukleinsäure nur dann hybridisiert, wenn die Sequenz des Oligonukleotids komplementär zu einer Region der amplifizierten Nukleinsäure ist, unter hybridisierenden Bedingungen mit dem in Schritt (c) erhaltenen Produkt behandelt wird; und

(e) nachgewiesen wird, ob eine amplifizierte Nukleinsäure in der Probe mit dem an der Membran befestigten Oligonukleotid hybridisiert hat.

11. Verfahren zum Nachweis des Vorhandenseins einer spezifischen Nukleotidsequenz in einer Nukleinsäure in einer Probe, bei welchem:

(a) die Probe gleichzeitig oder nacheinander mit vier verschiedenen Nukleosidtriphosphaten, einem Mittel zur Polymerisation der Nukleosidtriphosphate und zwei Oligonukleotidprimern für die Nukleinsäure unter hybridisierenden Bedingungen behandelt wird, so daß ein Primer an die Nukleinsäure hybridisiert und ein Extensionsprodukt des Primers synthetisiert wird, das komplementär zu der Nukleinsäure ist, wobei die Primer so ausgewählt sind, daß das mit einem Primer synthetisierte Extensionsprodukt als Matrize für die Synthese des Extensionsprodukts des anderen Primers dienen kann, wenn es von seinem Komplement getrennt wird;

(b) die Probe unter denaturierenden Bedingungen behandelt wird, um die Primerextensionsprodukte von ihren Matrizen abzutrennen;

(c) die Probe gleichzeitig oder nacheinander mit den vier Nukleosidtriphosphaten, einem Mittel zur Polymerisation der Nukleosidtriphosphate und Oligonukleotidprimern behandelt wird, so daß ein Primerextensionsprodukt unter Verwendung jeder der in Schritt (b) hergestellten Einzelstränge als Matrize synthetisiert wird, wobei die Schritte (b) und (c) ausreichend oft wiederholt werden, um exponentiell die Menge der Nukleinsäure zu erhöhen und zu einer nachweisbaren Amplifikation davon zu führen;

(d) an einer Membran eine markierte sequenzspezifische Oligonukleotidsonde befestigt wird, die mit der amplifizierten Nukleinsäure nur dann hybridisiert, wenn die Sequenz der Sonde komplementär zu einer Region der amplifizierten Nukleinsäure ist;

(e) die Membran aus (d) unter hybridisierenden Bedingungen mit dem in Schritt (c) erhaltenen Produkt behandelt wird;

(f) das Produkt aus Schritt (e) mit einem Restriktionsenzym behandelt wird, das jedes gebildete Hybrid spalten wird, wenn die Sonde und jede nachzuweisende Sequenz beide eine Restriktionsstelle enthalten, die von dem Enzym erkannt wird; und

(g) nachgewiesen wird, ob ein markiertes Restriktionsfragment der erforderlichen Länge in der Restriktionsverdauung vorhanden ist, wodurch angezeigt wird, daß Hybridisierung auftrat.

12. Verfahren nach Anspruch 10, wobei die Primer Oligodeoxyribonukleotide sind und gegebenenfalls markiert sind und das Mittel zur Polymerisation eine DNA-Polymerase ist.

13. Verfahren nach Anspruch 10 oder Anspruch 12, wobei die Nukleinsäure DNA ist, gegebenenfalls menschliche genomische DNA.

14. Verfahren nach Anspruch 10 oder Anspruch 12, wobei die Nukleinsäure RNA ist.

15. Verfahren nach Anspruch 13, wobei die Primer ein Segment eines Klasse II-HLA-Gens amplifizieren, gegebenenfalls ein DQ-alpha-Allel, ein DQ-beta-Allel, ein DP-alpha-Allel, ein DP-beta-Allel, ein DR-alpha-Allel oder ein DR-beta-Allel.

16. Kit zum Nachweis des Vorhandenseins einer spezifischen Nukleotidsequenz in einer Nukleinsäure in einer Probe, wobei das Kit, in verpackter Form, eine Mehrbehältereinheit aufweist, mit:

(a) zwei Oligonukleotidprimern für die Nukleinsäure, wobei die Primer so ausgewählt sind, daß das mit einem Primer synthetisierte Extensionsprodukt als Matrize für die Synthese des Extensionsprodukts des anderen Primers dienen kann, wenn es von seinem Komplement getrennt wird, um die Nukleinsäure exponentiell zu amplifizieren;

(b) einem Mittel zur Polymerisation;

(c) vier verschiedenen Nukleosidtriphosphaten; und entweder:

(d) einer sequenzspezifischen Oligonukleotidsonde, die mit der amplifizierten, direkt auf die Membran aufgetüpfelten, Nukleinsäure nur dann hybridisiert, wenn die Sequenz in der Sonde komplementär zu einer Region der amplifizierten Nukleinsäure ist; und

(e) einer Membran, an welche die amplifizierte Nukleinsäure zum anschließenden Nachweis durch Sondenhybridisierung direkt binden kann; oder

(f) einer an einer Membran befestigten sequenzspezifischen Oligonukleotidsonde, wobei die Sonde mit der amplifizierten Nukleinsäure nur dann hybridisiert, wenn die Sequenz der Sonde komplementär zu einer Region der amplifizierten Nukleinsäure ist.

17. Kit nach Anspruch 16, wobei entweder die Primer Oligodeoxyribonukleotide sind, die Nukleosidtriphosphate dATP, dCTP, dGTP und TTP sind und die Sonde markiert ist, oder wobei mindestens einer der Primer und/oder mindestens eines der Nukleosidtriphosphate markiert ist.

18. Kit nach Anspruch 16 oder Anspruch 17, welches außerdem ein Mittel zur Trennung der Stränge jeder in der Probe vorhandenen doppelsträngigen Nukleinsäure enthält und/oder eine positive Kontrolle für die Sonde, die eine oder mehrere Nukleinsäuren mit der nachzuweisenden spezifischen Nukleotidsequenz enthält, und eine negative Kontrolle für die Sonde, die keine Nukleinsäure mit der nachzuweisenden Nukleotidsequenz enthält.

19. Kit nach einem der Ansprüche 16 bis 18, wobei das Mittel zur Polymerisation ein Enzym ist, ausgewählt aus E. coli-DNA-Polymerase I, klenow-Fragment von E. coli-DNA-Polymerase I, T4-DNA-Polymerase, einem hitzestabilen DNA-Polymeraseenzym und einer reversen Transkriptase.

20. Kit nach einem der Ansprüche 16 bis 19, wobei die spezifische Nukleotidsequenz innerhalb einer menschlichen Nukleinsäure enthalten ist, ausgewählt aus einem beta-Globingen, einem HLA-DQ-alpha-, -DQ-beta-, DR-beta-, DP-alpha- und DP-beta-Gen, und mitochondrialer DNA, wobei gegebenenfalls die Nukleotidvariation in der Sequenz mit insulinabhängigem Diabetes mellitus assoziiert ist.

## Revendications

1. Procédé de détection de la présence d'une séquence de nucléotides spécifique dans un échantillon d'acide nucléique, selon lequel :

(a) on traite l'échantillon dans des conditions d'hybridation, simultanément ou successivement, avec quatre nucléoside triphosphates différents, un agent de polymérisation des nucléoside triphosphates, et deux amorces oligonucléotidiques correspondant à l'acide nucléique, de sorte qu'une amorce s'hybride avec l'acide nucléique, et un produit d'extension de l'amorce complémentaire de l'acide nucléique, est synthétisé, les amorces étant choisies de telle sorte que le produit d'extension synthétisé à partir d'une amorce, puisse, lorsqu'il est séparé de son complément, servir de matrice pour la synthèse du produit d'extension de l'autre amorce ;

(b) on traite l'échantillon dans des conditions de dénaturation afin de séparer les produits d'extension d'amorce

de leurs matrices ;

(c) on traite l'échantillon, simultanément ou successivement, avec les quatre nucléoside triphosphates, un agent de polymérisation des nucléoside triphosphates, et des amorces oligonucléotidiques, de telle sorte qu'un produit d'extension d'amorce est synthétisé en employant chacun des simples brins produits dans l'étape (b) comme matrice, les étapes (b) et (c) étant répétées un nombre de fois suffisant pour accroître exponentiellement la quantité de l'acide nucléique, et obtenir une amplification détectable de celui-ci ;

(d) on transfère directement sur une membrane, sans fractionnement sur gel, le produit issu de l'étape (c) ;

(e) on traite la membrane obtenue en (d) dans des conditions d'hybridation, avec une sonde oligonucléotidique marquée spécifique d'une séquence, capable de s'hybrider avec l'acide nucléique amplifié, seulement si la séquence de la sonde est complémentaire d'une région de l'acide nucléique amplifié ; et

(f) on détecte si la sonde a été hybridée avec un acide nucléique amplifié présent dans l'échantillon.

2. Procédé selon la revendication 1, dans lequel l'acide nucléique présent dans l'échantillon, est extrait de l'échantillon avant l'étape (a), et dans lequel on répète les étapes (b) et (c) au moins cinq fois, et éventuellement dans le cas où l'échantillon contient de l'ADN de génome humain, les étapes (b) et (c) sont répétées 15 à 30 fois.

3. Procédé selon la revendication 1 ou 2, dans lequel l'acide nucléique est bicaténaire, et ses brins sont séparés par dénaturation avant ou au cours de l'étape (a).

4. Procédé selon la revendication 1 ou 2, dans lequel l'acide nucléique est monocaténaire.

5. Procédé selon la revendication 3, dans lequel l'acide nucléique est un ADN, et les amorces sont des oligodésoxyribonucléotides.

6. Procédé selon la revendication 4, dans lequel l'acide nucléique est un ADN ou un ARN messager, et les amorces sont des oligodésoxyribonucléotides.

7. Procédé selon l'une quelconque des revendications 1 à 6, dans lequel l'agent de polymérisation est choisi parmi l'ADN polymérase I de *E. coli*, le fragment de Klenow de l'ADN polymérase I de *E. coli*, l'ADN polymérase T4, une polymérase thermostable et une transcriptase reverse.

8. Procédé selon l'une quelconque des revendications 1 à 7, dans lequel la séquence de nucléotides spécifique, est contenue dans un acide nucléique humain choisi parmi un gène de  $\beta$ -globine, un oncogène, un gène HLA-DQ alpha, un gène DQ bêta, un gène DR bêta, un gène DP alpha, un gène DP bêta et un ADN mitochondrial, et dans lequel la séquence de nucléotides est éventuellement associée au diabète sucré dépendant de l'insuline, ou dans lequel la séquence de nucléotides est contenue dans l'acide nucléique d'un organisme qui peut provoquer une maladie infectieuse.

9. Procédé selon l'une quelconque des revendications 1 à 8, dans lequel dans le cas où les échantillons comprennent des cellules, éventuellement lorsque l'échantillon comprend des lymphocytes de sang périphérique ou un liquide amniotique, le procédé comprend en outre avant l'étape (a), l'étape de chauffage de l'échantillon dans une mesure suffisante pour exposer le ou les acides nucléiques contenus dans celui-ci, ou dans le cas où l'échantillon est un échantillon de sperme, de cheveux ou de sang, ou s'il contient de l'ADN mitochondrial, le procédé est employé pour l'analyse judiciaire.

10. Procédé de détection de la présence d'une séquence de nucléotides spécifique dans un échantillon d'acide nucléique, selon lequel :

(a) on traite l'échantillon dans des conditions d'hybridation, simultanément ou successivement, avec quatre nucléoside triphosphates différents, un agent de polymérisation des nucléoside triphosphates et deux amorces oligonucléotidiques correspondant à l'acide nucléique, de telle sorte qu'une amorce s'hybride avec l'acide nucléique, et un produit d'extension de l'amorce complémentaire de l'acide nucléique, est synthétisé, les amorces étant choisies de telle sorte que le produit d'extension synthétisé à partir d'une amorce puisse, lorsqu'il est séparé de son complément, servir de matrice pour la synthèse du produit d'extension de l'autre amorce ;

(b) on traite l'échantillon dans des conditions de dénaturation, pour séparer les produits d'extension d'amorce de leurs matrices ;

(c) on traite l'échantillon, simultanément ou successivement, avec les quatre nucléoside triphosphates, un agent de polymérisation des nucléoside triphosphates, et des amorces oligonucléotidiques, de telle sorte qu'un

produit d'extension d'amorce soit synthétisé en employant comme matrice, chacun des simples brins produits dans l'étape (b), les étapes (b) et (c) étant répétées un nombre de fois suffisant pour accroître exponentiellement la quantité de l'acide nucléique, et obtenir un degré d'amplification détectable de celui-ci ; et dans lequel on marque au moins une amorce et/ou au moins l'un des quatre nucléoside triphosphates, avec un groupe détectable ;

(d) on traite dans des conditions d'hybridation avec le produit de l'étape (c), une membrane à laquelle est lié un oligonucléotide spécifique d'une séquence capable de s'hybrider avec l'acide nucléique amplifié, seulement si la séquence de l'oligonucléotide est complémentaire d'une région de l'acide nucléique amplifié ; et

(e) on détecte si un acide nucléique amplifié présent dans l'échantillon, s'est hybridé avec l'oligonucléotide lié à la membrane.

11. Procédé de détection de la présence d'une séquence de nucléotides spécifique dans un échantillon d'acide nucléique, selon lequel :

(a) on traite l'échantillon dans des conditions d'hybridation, simultanément ou successivement, avec quatre nucléoside triphosphates différents, un agent de polymérisation des nucléoside triphosphates, et deux amorces oligonucléotidiques correspondant à l'acide nucléique, de telle sorte qu'une amorce s'hybride avec l'acide nucléique, et un produit d'extension de l'amorce complémentaire de l'acide nucléique, est synthétisé, les amorces étant choisies de telle sorte que le produit d'extension synthétisé à partir d'une amorce, puisse, lorsqu'il est séparé de son complément, servir de matrice pour la synthèse du produit d'extension de l'autre amorce ;

(b) on traite l'échantillon dans des conditions de dénaturation, pour séparer les produits d'extension d'amorce de leurs matrices ;

(c) on traite l'échantillon, simultanément ou successivement, avec les quatre nucléoside triphosphates, un agent de polymérisation des nucléoside triphosphates, et les amorces oligonucléotidiques, de telle sorte qu'un produit d'extension d'amorce soit synthétisé en employant comme matrice, chacun des simples brins produits dans l'étape (b), les étapes (b) et (c) étant répétées un nombre de fois suffisant pour accroître de manière exponentielle la quantité de l'acide nucléique, et obtenir un degré d'amplification détectable de celui-ci ;

(d) on lie à une membrane, une sonde oligonucléotidique marquée spécifique d'une séquence, capable de s'hybrider avec l'acide nucléique amplifié, seulement si la séquence de la sonde est complémentaire d'une région de l'acide nucléique amplifié ; et

(e) on traite dans des conditions d'hybridation, la membrane obtenue en (b), avec le produit issu de l'étape (c) ;

(f) on traite le produit de l'étape (e) avec une enzyme de restriction qui coupe les hybrides formés si la sonde et une séquence quelconque destinée à être détectée, contiennent toutes les deux un site de restriction reconnu par l'enzyme ; et

(g) on détecte si un fragment de restriction marqué ayant la longueur requise, est présent dans le produit de digestion par restriction, ce qui indique que l'hybridation a eu lieu.

12. Procédé selon la revendication 10, dans lequel les amorces sont des oligodésoxyribonucléotides éventuellement marqués, et dans lequel l'agent de polymérisation est une ADN polymérase.

13. Procédé selon la revendication 10 ou 12, dans lequel l'acide nucléique est un ADN, éventuellement un ADN de génome humain.

14. Procédé selon la revendication 10 ou 12, dans lequel l'acide nucléique est un ARN.

15. Procédé selon la revendication 13, dans lequel les amorces amplifient un segment d'un gène HLA de classe II, éventuellement un allèle DQ-alpha, un allèle DQ-béta, un allèle DP-alpha, un allèle DP-béta, un allèle DR alpha ou un allèle DR bêta.

16. Nécessaire pour la détection de la présence d'une séquence de nucléotides spécifique dans un échantillon d'acide nucléique, comprenant, dans un emballage, un ensemble de récipients comportant :

(a) deux amorces oligonucléotidiques correspondant à l'acide nucléique, ces amorces étant choisies de telle sorte qu'un produit d'extension synthétisé à partir d'une amorce, puisse, lorsqu'il est séparé de son complément, servir de matrice pour la synthèse du produit d'extension de l'autre amorce, de façon à amplifier exponentiellement l'acide nucléique ;

(b) un agent de polymérisation ;

(c) quatre nucléoside triphosphates différents ; et

soit :

(d) une sonde oligonucléotidique spécifique d'une séquence, capable de s'hybrider avec l'acide nucléique amplifié déposé directement sur une membrane seulement si la séquence de la sonde est complémentaire d'une région de l'acide nucléique amplifié; et

(e) une membrane à laquelle l'acide nucléique amplifié peut se lier directement afin de permettre une détection ultérieure par hybridation avec une sonde ; soit

(d) une séquence oligonucléotidique spécifique d'une séquence liée à une membrane, la sonde étant capable de s'hybrider avec l'acide nucléique amplifié, seulement si la séquence de la sonde est complémentaire d'une région de l'acide nucléique amplifié.

17. Nécessaire selon la revendication 16, chaque amorce étant un oligodésoxyribonucléotide, les nucléoside triphosphates comprenant le dATP, le dCTP, le dGTP et le TTP, et la sonde étant marquée ; ou dans lequel au moins l'une des amorces et/ou au moins l'un des nucléoside triphosphates, sont marqués.

18. Nécessaire selon la revendication 16 ou 17, comprenant en outre des moyens de séparation des brins des acides nucléiques bicaténaire contenus dans l'échantillon, et/ou un témoin positif pour la sonde, contenant un ou plusieurs acides nucléiques avec la séquence nucléotidique spécifique destinée à être détectée, et un témoin négatif pour la sonde, ne contenant pas d'acide nucléique avec la séquence nucléotidique destinée à être détectée.

19. Nécessaire selon l'une quelconque des revendications 16 à 18, dans lequel l'agent de polymérisation est une enzyme choisie parmi l'ADN polymérase I de E. coli, le fragment de Klenow de l'ADN polymérase I de E. coli, l'ADN polymérase T4, une ADN polymérase thermostable, et une transcriptase reverse.

20. Nécessaire selon l'une quelconque des revendications 16 à 19, dans lequel la séquence de nucléotides spécifique, est contenue dans un acide nucléique humain choisi parmi un gène de bêta-globine, un gène HLA-DQ alpha, un gène DQ bêta, un gène DR bêta, un gène DP alpha, un gène DP bêta et un ADN mitochondrial, la variation de nucléotide dans la séquence étant associée au diabète sucré dépendant de l'insuline.